



Bayer Crop Science

Assessment of on-field GHG emission of crops

Draft report – 3rd review cycle

December 2022

Note to reviewers:

Through this critical review, Bayer aims to verify that it uses the Cool Farm Tool (CFT) for greenhouse gas (GHG) emission quantification in a reasonable approach and that the baselining and performance tracking methodology is adequate.

Disclaimer

This material may contain “forward-looking statements” based on current assumptions and forecasts made by Bayer management. Various known and unknown risks, uncertainties and other factors could lead to material differences between the actual future results, financial situation, development or performance of the company and the estimates given here. These factors include those discussed in Bayer’s public reports which are available on the Bayer website at <http://www.bayer.com>. The company assumes no liability whatsoever to update these forward-looking statements or to conform them to future events or developments.

Contents

1	Context and Objectives.....	6
1.1	Context.....	6
1.2	Review of GHG emissions related to agriculture, forestry and land use activities with BCS' role in agriculture-linked GHG reduction.....	7
1.3	Bayer GHG reduction target consistent with its commitment to international frameworks and key initiatives.....	8
1.4	Objectives.....	9
1.5	Critical review	9
1.6	Organization of the study.....	10
1.7	Use of the study and target audience	11
2	Scope	11
2.1	Aggregated system studied: From individual farms to crop-country combinations (CCCs) and rationale for their selection	11
2.2	System Boundaries: Defining the scope of the estimated emissions.....	12
2.3	Functional unit	13
3	Method	13
3.1	Description of the GHG Assessment Inventory data	13
3.1.1	Sampling approach and processing	14
3.1.2	Data quality check by Kynetec.....	15
3.1.3	Inventory data compilation for the GHG assessment	15
3.1.4	CCCs production quantity and BCS market share.....	16
3.1.5	Definition of BCS customer base used for the on-field GHG assessment	17
3.2	Determination of on-field GHG emissions and carbon sequestration with the Cool Farm Tool 18	
3.2.1	Cool Farm Tool model description	18
3.2.2	Cool Farm Tool input data	19
3.3	Calculation of BCS Customers GHG emission.....	26
3.3.1	Calculation of specific GHG baseline for particular CCC's.....	26
3.3.2	Setting an aggregated baseline for GHG emission reduction across CCCs	27
4	Interpretation	28
4.1	Results and setting of GHG emission tracking baseline.	28
4.1.1	Specific GHG emissions for the baseline	29
4.1.2	Total GHG emissions across the CCC for the baseline	30
4.1.3	Aggregated GHG emission for the baseline year.....	31
4.1.4	Performance tracking	32
4.2	Discussion	33
4.3	Uncertainty analysis discussion in extant literature.....	34
4.4	Sensitivity analysis discussion in extant literature	34

5	Main limitations of the assessment	36
6	Further developments of this report	36
7.	References	36
8.	Appendix.....	39

List of Figures

Figure 1 - Main sources and sinks of emissions from agricultural systems. Figure taken from IPCC (2006)	7
Figure 2 - Workflow for the calculation of farm level carbon footprints	14
Figure 3 - Qualifying criteria to be met by farmers for selection as part of survey respondents.....	14
Figure 4 - A stratified sampling plan for data collection on Maize (same as corn) cultivation in Belgium. .	15
Figure 5 - Specific GHG emission in the baseline based on BCS'S customer farms.....	30
Figure 6 - Specific GHG emission in the baseline based on BCS'S customer farms grouped by emission sources	30
Figure 7 - Total GHG emission based on BCS's customer farms for the 18 CCC's	31
Figure 8 - Individual weighting factors for the 18 CCCs	32
Figure 9 - Baseline establishment and performance tracking	33
Figure 10 - Factors that contribute to the variability in the carbon footprint results from CFT GHG calculation (Figure from Clavreul et al. (2017)).....	35

List of Tables

Table 1 - Critical review panel composition.....	10
Table 2 - Contact information for all parties	11
Table 3 - Overview of activities included in the system boundaries	12
Table 4 - Kynetec data collection phases and total sample size for the crop-country-combination's	16
Table 5 - Summary of UNFAO reported average crop production for 2015-2019 and BCS market share for the CCCs.....	16
Table 6 - Sample size and number of customers according to definition of BCS's customer base.	18
Table 7 - Cool Farm Tool Input parameters on farm settings	20
Table 8 - Cool Farm Tool Input parameters on crop details	20
Table 9 - Cool Farm Tool Input parameters on crop residue management	21
Table 10 - Cool Farm Tool Input parameters on soil characteristics.....	21
Table 11 - Cool Farm Tool Input parameters on fertilizer management	22
Table 12 - Cool Farm Tool Input parameters on direct energy use.....	23
Table 13 - Cool Farm Tool Input parameters on cultivation practices and field operations.....	24
Table 14 - Cool Farm Tool Input parameters on field operations energy use.....	24
Table 15 - Cool Farm Tool Input parameters on tillage and cover crops management.....	25
Table 16 - Specific GHG emission (kg CO ₂ e per kg crop) from surveyed BCS customer farms in the baseline	29
Table 17 - Total GHG emissions (kg CO ₂ e) in the baseline year and different factors used in calculating Bayer's total GHG emissions in the 18 CCCs.	31
Table 18 - The specific GHG emission, total GHG emission and weighting factors for the 18 CCCs	32
Table 19 - Summary of specific GHG emission (kg CO ₂ e per kg crop) from surveyed BCS customer farms by gate-to-gate emission source in the baseline year for the 18 CCC's.	39

Table of Abbreviations

ANA Brazil	Agência Nacional de Águas e Saneamento Básico, Brasil National Water and Basic Sanitation Agency, Brazil
BCS	Bayer Crop Science
CCCs	Crop-country combinations
CFT	Cool Farm Tool
CH ₄	Methane
CO ₂	Carbon dioxide
CO ₂ -eq	Carbon dioxide equivalent
CPP	Crop protection product
EASAC	European Academies Science Advisory Council
FAO	Food and Agriculture Organization of the United Nations
FAQ	Frequently asked questions
GHG	Greenhouse gas
Gt	Gigatons
ha	Hectare
ILCD	International Reference Life Cycle Data System
IPCC	Intergovernmental Panel on Climate Change
kg	Kilogram
kWh	Kilowatt per hour
LCA	Life Cycle Assessment
N	Nitrogen
N ₂ O	Nitrous oxide
NH ₃	Ammonia
NO	Nitric oxide
pH	Potential of Hydrogen - Logarithmic expression of the hydrogen ions concentration in a solution
SBTi	Science Based Targets initiative
SDGs	Sustainable Development Goals
t	Ton
UK	United Kingdom
UN	United Nations
US	United States of America
USDA-NASS	U.S. Department of Agriculture - National Agricultural Statistics Service
WIR	World Resources Institute
WWF	World Wide Fund for Nature
yr	Year

1. Context and Objectives

1.1 Context

Bayer is a Life Science company with a more than 150-year history and core competencies in the areas of health care and nutrition. Contributing to sustainable development has become a core element of Bayer's corporate strategy. For Bayer Crop Science (BCS) division, sustainability focus areas and goals were developed to fulfill the commitment to shape the future of sustainable agriculture. BCS' sustainability focus areas were developed to address the end-to-end impacts of agriculture, which include transformational commitments on field greenhouse gas (GHG) emissions, on environmental impact reduction of crop protection, improving the livelihoods of smallholder farmers. Further focus areas address biodiversity & soil health, product responsibility (ensuring the safe usage of crop protection products) as well as conserving & protecting water resources.

According to the Intergovernmental Panel on Climate Change (IPCC), the GHG emissions from the global food system are estimated to be 21-37% of total net anthropogenic GHG emissions (IPCC, 2019). As one of the largest agricultural companies in the world, Bayer recognizes the impact of its products and aims to empower farmers to reduce the on-field GHG emissions of agriculture wherever the company operates. As part of its sustainability objectives, BCS has committed to a **30% reduction of on-field GHG emissions per kilogram of crop produced by BCS's farming customers by 2030 for the most emitting crop systems in the regions where BCS operates (i.e., the BCS on-field GHG commitment)**. The scope of BCS' efforts is focused on emissions of major GHGs (CO₂, CH₄, N₂O) from the field operations. To meet this objective, Bayer aims to foster and encourage the adoption of climate-smart practices and technologies amongst its farming customer base.

The main objective of this report is to document how BCS is quantifying field GHG emissions and soil carbon sequestration. More specifically, this report documents how BCS compiles inventory data and conducts a GHG impact assessment based on the internationally recognized and empirically validated Cool Farm Tool (CFT) calculator¹. The CFT will further be used in the determination of improvement potentials towards the GHG reduction target. While being aware of the potential risk of burden shifting, BCS emphasizes that this assessment focuses on the GHG emissions and soil carbon sequestration resulting from field operations and does not cover other impact categories such as ecotoxicity and other [BCS sustainability focus areas](#) as they are assessed and documented in separate reports by different task forces.

In addition to setting targets on the GHG resulting from farming, BCS has committed to enable farmers to reduce the ecotoxicity impact from crop protection use by 30%, by the year 2030. BCS also strives to improve the livelihoods of 100 million smallholder farmers through access to education and tailored solutions. In addition, strategies are being developed towards biodiversity and soil health, water conservation and product responsibility for the promotion of sustainability in agriculture. In the context of this report, BCS does not conduct a full-fledged LCA according to ISO 14040/44 but intends to use the standard as a framework to document the project in the present report. With the critical review by external experts, BCS aims to verify that it uses the CFT calculator in a reasonable approach and that the baselining and performance tracking methodology is adequate. In case of external communication of the present report or any material based on it, BCS intends to publish the external expert panels feedback with transparency, and it intends to consult the panel regularly in the future.

¹ The cool farm tool (CFT) developed by the Cool Farm Alliance is a tool used to measure greenhouse gas emissions from agricultural production of a specific product and on a global scale. See section 0 for more details.

1.2 Review of GHG emissions related to agriculture, forestry and land use activities with BCS' role in agriculture-linked GHG reduction

Food related emissions are those generated during production activities (crops and livestock), land use change and pre- and post-production processes. Production and land use change result in emissions generated on agricultural land, while pre- and post-production refer to emissions from supply chain processes including transportation, processing and manufacturing of inputs. In 2019, the global anthropogenic emissions were estimated to be 54 billion tonnes of CO₂eq in which 17 billion tonnes CO₂eq (31%) comes from agricultural related activities. Breaking the share of agricultural related sources (31%) from the total anthropogenic emissions down to single gases, CO₂ accounts for 21%, methane (CH₄) accounts for 53% while nitrous oxide (N₂O) accounts for the highest which is 78 % (FAO., 2021). Aligning current production and consumption models in the agri-food sector with planetary boundaries² is vital for constructing a resilient food system and ensuring companies continue to thrive in a resource-constrained world.

According to the FAO, (2021), farmgate emissions account for the largest share of the agricultural related emissions in 2019 with about 7 billion tonnes CO₂eq. While agriculture plays a role in GHG emission (Figure 1), climate change on the other hand also places significant pressures on agriculture in the form of reduced yields, land degradation, and increased threats from pathogens and disease. That means agriculture is confronted with tremendous challenges regarding climate change mitigation and adaptation.




Figure 1 - Main sources and sinks of emissions from agricultural systems. Figure taken from IPCC (2006)
CH₄: methane, CO₂: carbon dioxide, N₂O: nitrous dioxide, NO_x: Nitrogen oxides, CO: Carbon monoxide, NMVOC: non-methane volatile organic compounds, HWP: harvested wood products.

BCS has a responsibility to advance a net zero future for agriculture. Great progress has already been made to reduce agriculture's overall carbon footprint, but BCS must work collectively with farmers and global partners to do even more. These requires innovation and huge advancements in agricultural

² As defined by Steffen, 2015, "The planetary boundaries framework defines a safe operating space for humanity based on the intrinsic biophysical processes that regulate the stability of the Earth system – Steffen et al., by (2015) "

technologies. To accelerate this shift, Bayer developed ambitious and robust targets building on the state-of-the-art and internationally recognized CFT calculator.

1.3 Bayer GHG reduction target consistent with its commitment to international frameworks and key initiatives

Bayer AG (including the BCS division) is part of the world's leading Science Based Targets initiative (SBTi), a joint initiative of the Carbon Disclosure Project (CDP), the United Nations Global Compact, the World Resources Institute (WRI) and the World Wide Fund for Nature (WWF). SBTi focuses on providing companies with a scientifically-based framework for setting ambitious and effective climate targets towards the long-term goal of achieving net-zero emissions. It outlines criteria for effective reduction of companies' greenhouse gas (GHG) emissions in line with the Paris Agreement goal of limiting global warming to 1.5 °C, compared to pre-industrial levels. SBTi further validates set corporate goals to ascertain that they are in line with its criteria.

The Bayer AG has the aim to continuously reduce GHG emissions within the company and along the entire value chain in accordance with the set criteria and validation of the SBTi. In line with this, Bayer has set a target to achieve net zero GHG emissions including its entire value chain by 2050 or sooner. By 2024 (interim target), Bayer plans to reduce its scope 1&2³ emissions by 20% and its Scope 3 emissions by 6% with reference to the baseline year 2019. Until the end of 2029 (Mid-term target), it plans to reduce its scope 1 & 2 emissions by 42% and its scope 3 emission through cooperation with suppliers and customers by at least 12.3% compared to its 2019 baseline. To accomplish this, Bayer will combine measures, such as more efficient inward and outward ventilation systems, a move to climate-neutral technologies, such as geothermal energy for heating and cooling and a switch to 100% purchased electricity from renewable sources. These targets have been approved by the Science Based Target initiative as aligned with a 1.5°C pathway for Scope 1 and 2 and with a 2°C pathway for Scope 3 emissions. As such targets cannot be achieved by acting alone, Bayer has joined forces with other ambitious companies to drive progress as a part of the chemical industry's "Together for Sustainability" initiative. Bayer is also a member of the CDP Supply Chain Initiative and in direct contact with key suppliers.

Additionally, Bayer is on a path to become climate neutral by 2030 in its own operations. The remaining emissions after reduction will be offset by purchasing certificates from climate protection projects with recognized quality standards. The offset projects are related to our business. Based on our business purpose we focus on Natural Climate Solutions relating to forest and agriculture. Additionally, we invest in innovative projects and foster development of voluntary carbon markets.

The BCS division is also planning additional climate protection measures that go beyond the Bayer-AG-wide GHG reduction targets outlined above. With the BCS GHG commitment, BCS aims to enable its farming customers to reduce their field crops emissions per kg of crop produced by 30% in 2030.

Therefore, BCS will contribute to reduction of on-field GHG emissions and promote soil carbon sequestration among its customers in relevant crops and geographies. BCS will do so by leveraging their expertise and innovative seeds and crop protection portfolio, promoting the use of modern and efficient farming practices as well as capitalizing on its digital farming solutions. Together with its partners, BCS will strive to promote climate-smart solutions and combine different levers to profitable/customized tailored solutions that help farmers to increase their resilience to consequences of climate change (such as droughts, heavy rains, erosion). Consequently, BCS will bolster farmers with the right tools and technologies to sequester carbon in the soil, reduce and avoid emissions and grow crops in a sustainable manner. Such levers include high yielding crop genetics, crop protection agents, water use efficiency, soil management through no-till and cover crops, crop rotation, root health, (nitrogen-) fertilization management, shortening

³ Scope 1 emissions are direct GHG emissions that occur from company owned or controlled sources (e.g., emissions associated with fuel combustion in boilers, vehicles etc.), Scope 2 emissions are indirect GHG emissions associated with the purchase of electricity, steam, heat, or cooling while Scope 3 emissions are all indirect emissions (not included in scope 2) that occur in the value chain.

the time of flooding in rice (direct seeded rice), digital tools to support decision processes and use of biological CPP (biologics).

Thus, the BCS on-field GHG reduction commitment will contribute to attaining the United Nations' Sustainable Development Goals (UN SDGs). The United Nations agreed on 17 SDGs to build a better world for people and our planet by 2030. The 2030 Sustainable Development Agenda emphasizes that development should be compatible with all three dimensions of sustainability: economic, social, and environmental. Implementing the 2030 Agenda presents an opportunity for collaborative action by many diverse actors, and at all levels, to minimize adverse climate change impacts of agriculture. Therefore, BCS' on-field GHG commitment is at the interface with several goals of the 2030 Agenda (UN, United Nations, 2021) to contribute to sustainable farming practices and food production:

- SDG 2 – End hunger, achieve food security and improved nutrition and promote sustainable agriculture. Considering that the on-field GHG commitment also builds on yield improvements, BCS will specifically contribute to the SDG targets 2.1 (end hunger), 2.3 (increase the agricultural productivity), and 2.4 (ensure sustainable food production systems and implement resilient agricultural practices that increase productivity and production, that help maintain eco-systems, that strengthen capacity for adaptation to climate change, extreme weather, drought, flooding and other disasters and that progressively improve land and soil quality).
- SDG 13 – Take urgent action to combat climate change and its impacts.

1.4 Objectives

In 2019, Bayer publicly committed to equipping farmers to reduce the agricultural GHG emissions per kilogram of crop produced by 30% by the year 2030 with a focus on where its products are used. To deliver on this goal, the first step is to establish a baseline against which to measure progress. BCS has set the foundations for its performance tracking method based on CFT to understand the current on-field GHG emissions. Establishing a baseline will help identify opportunities for reducing GHG emissions.

Therefore, this report is aimed at achieving the below objectives:

- Document a method to quantify BCS farming customers' on-field GHG emissions and soil carbon sequestration in order to account for the climate change impacts resulting from farming operations on the field.
- Determine improvement potentials in line with the Bayer on-field GHG reduction target.

To achieve this, we will be using the CFT GHG emission quantification tool and inventory data from Kynetec to account for the GHG emissions from BCS farming customers'. Based on this method, BCS will calculate a baseline in order to track performance and progress against the 30% on-field GHG reduction commitment.

1.5 Critical review

This report is structured using the Life Cycle Assessment (LCA) methodology (according to the ISO 14040 and ISO 14044) as a template for documentation of methodological choices, results, and interpretations as well as limitations. As such, BCS acknowledges that this report only focuses on the field gate-to-gate⁴ life cycle stage GHG emissions and soil carbon sequestration resulting from farming operations. Consequently, BCS does not claim that this report complies with ISO 14040/44. As BCS intends to communicate to the public its sustainability commitments and achievements, a critical review has been

⁴ Field gate-to-gate refers to the GHG emission resulting from crop production, starting from on-field soil preparation until the moment the crop leaves the farmers' field.

performed, following a three-step iterative process. This report provides the review panel composition, its conclusions and the details of the comments and final report adaptations.

Table 1 - Critical review panel composition

Members	Country	Area of expertise
Thomas Nemecek	Switzerland	Deputy Lead Life Cycle Assessment Research Group Agroscope. Worldwide known researcher on Life Cycle Assessment, specifically in its applications on agriculture.
Jeffrey Jenkins	U.S.A.	Expertise in environmental analytical chemistry, ecological risk assessment, and agronomically-based ecohydrologic modeling to characterize watershed scale pesticide use and the potential impact on water quality.
Valery Forbes	U.S.A.	Dean and Professor at Florida Atlantic University. Broad expertise in mechanistic effect modeling and ecological risk assessment of pesticides and other chemicals.
Assumpció Anton	Spain	Researcher at Food and Agricultural Research Institute, IRTA. Expertise in the development and application of LCA methodology in agriculture.
Tiago Rocha	Brazil	Consultant Partner at ACV Brasil and PhD in Environmental Technology. Extensive experience in life cycle assessment, specifically in the area of carbon footprint.
Lorie Hamelin	France	Researcher at the Federal University of Toulouse (France), studying the environmental impacts related to large-scale transitions towards low fossil carbon use.
Anne-Marie Boulay	Canada	Assistant Professor in Chemical Engineering at Polytechnique Montreal and CIRAIG. Expertise on water footprint methodologies and impact assessment associated with plastic litter in LCA.
Jessica Hanafi	Indonesia	PhD in Life Cycle Engineering. Established the Indonesian Association of Life Cycle Assessment and Sustainability Professional. ISO Technical Committee on Life Cycle Assessment (TC 207/SC5), environmental labelling (SC3), Greenhouse Gas (SC7) and project leader for ISO/TS 14074 LCA normalization and weighting. Applied LCA based on ISO 14040/44 to various industrial sectors, including agriculture.
Laura Golsteijn (Chair of the panel)	Netherlands	Senior LCA Consultant at PRé. PhD in Toxic Impact Modelling. Supporting clients to understand, develop and embed environmental metrics to improve the sustainability of supply chains and products.

1.6 Organization of the study

The overall primary data collection and GHG impact calculation process can be summarized as follows: For the compilation of inventory data, BCS uses inventory data from Kynetec's 'FarmTrak™' which tracks global agriculture in 52 countries, by surveying and interviewing global grower panels annually and collecting details of the crops grown (Kynetec, 2021). These data are supplemented with FarmTrak Sustainability data which contain other field operation data like machinery and cultivation techniques. The combined data set compiles all relevant information related to seed, crop protection, fertilizer use, and yield. Based on these extensive crop input data sets, Kynetec calculates on-field GHG emissions following the calculation methodology of the Cool Farm Tool. Then, BCS interprets the results to set a global on-field GHG baseline value across Crop-Country Combinations (CCCs) and to determine improvement potentials. More details on the compilation of inventory data, impact assessment, and interpretation follow in later sections of this report.

Table 2 - Contact information for all parties

Organization	Task	Contact information (Role)
Bayer Crop Science	<ul style="list-style-type: none"> • Calculate global on-field GHG baseline value across CCCs. • Apply global on-field GHG baseline internally at BCS to determine improvement potentials in line with the Bayer on-field GHG reduction commitment. • Assess how to integrate learnings into business models. Enable BCS organization to work on on-field GHG data. 	Dr. Alexey Kuzmenkin alexey.kuzmenkin@bayer.com (Director, Global Climate Change Lead)
Kynetec	<ul style="list-style-type: none"> • Data collection (based on FarmTrak™) and on-field GHG impact calculation (based on Cool Farm Tool procedures). 	Christophe Labyt christophe.labyt@kynetec.com (Director - Sustainability Products and Services at Kynetec) Stephen Hearn stephen.hearn@kynetec.com (Consultant to Leadership teams at Kynetec)

1.7 Use of the study and target audience

The results of this report are intended to transparently and publicly describe the baseline, performance tracking and GHG calculation method. BCS aims to publish the expert panels feedback as well to ensure transparency and strive for credibility. Therefore, the main target audience are investors, press, academic partners and the general public. Potentially, this report might also be used in the future for auditing processes.

This report is not BCS's main vehicle for informing external stakeholders. BCS is currently developing other internal and external training and communication materials and channels that will be specifically tailored to the information needs of the respective stakeholder group.

2 Scope

This section includes a description of the system boundaries, functional unit, and other relevant scenario and scope information.

2.1 Aggregated system studied: From individual farms to crop-country combinations (CCCs) and rationale for their selection

This report focuses on quantifying on-field GHG emissions and soil carbon sequestration to account for the most emitting crop systems in the regions where BCS operates. To achieve this, field-level GHG emission and sequestration results are aggregated to a higher level of crops and countries (so-called crop-country combinations; CCCs). Based on interview data with farmers from these 18 CCCs, BCS is quantifying the

climate change impacts of its farming customers within a whole CCC. The following 18 CCCs were selected for the assessment:

- Argentina-corn
- Argentina-soybeans
- Australia-cotton
- Australia-wheat
- Brazil-corn
- Brazil-soybeans
- Canada-rapeseed
- Canada-wheat
- France-wheat
- India-rice, paddy
- Italy-corn
- Mexico-corn
- Spain-corn
- USA-cotton
- USA-corn
- USA-soybeans
- USA-wheat (Spring)*⁵
- USA-wheat (Winter)*

The 18 CCCs were pre-determined and selected, with the intention of representing BCS's market share and influence for improvement. This allows BCS to target the crops where it has the largest potential for reduction in order to meet its sustainability-related objectives. The CCCs were selected based on the following criteria:

- Business relevance based on production volume of a particular crop in a particular market (FAO database) and Bayer market share in a particular market.
- Climate change relevance based on carbon footprint of the cropping systems and GHG reduction potential.
- Italy-Corn and Spain-Corn were not selected based on these factors, but were additionally included because data were already available.

2.2 System Boundaries: Defining the scope of the estimated emissions

This section provides an overview of the emissions included (in-scope) in this assessment. The specific GHG emission is determined within the gate-to-gate GHG emission from the survey of BCS farming customers based on the CFT methodology. The assessment excludes some emission categories that occurs beyond the farmers field and are such considered out-of scope. The assessment focuses on emissions that farmers can directly influence. Information on the emissions considered (in-scope) in this assessment are listed in Table 3 below.

Table 3 - Overview of activities included in the system boundaries

In-scope emissions	Details
Fertilizer application	includes on-field emissions from fertilizer decomposition, encompassing CO ₂ , N ₂ O, NO and NH ₃ emissions.
Energy sources consumed on the farm	includes farm machinery use during sowing, cultivation, application of fertilizer and crop protection products, harvesting, and irrigation.

⁵ *USA wheat is divided into spring and winter as a result of the differences in harvest season, management (fertilizer and crop protection) and yield.

Organic matter application	includes on-field emissions coming from decomposition of left-over residues, or from other ways of managing residue (incorporating it in the soil, taking it off field etc.).
Management changes	Includes changes in soil carbon stock due to soil management (tillage practices and cover crops), soil organic carbon accumulation or decline and carbon sequestration (a negative CO ₂ emission).

Although the CFT calculates emissions related to the production of crop protection products and fertilizers, transportation, drying and land use change, these emissions are considered out of scope in this assessment. Transportation is excluded because it refers to activities outside the farmgate and land use change is considered out of scope due to lack of reliable data. Drying activity is carried out off field, therefore excluded as the analysis was done to calculate emissions from sources that are within the farm-gate to farm-gate boundaries. The production of crop protection products and fertilizers is out of scope because of BCS's strategic decision to focus on on-field GHG emissions that farmers can directly influence.

2.3 Functional unit

Since the function of the system is to produce crop biomass for food, feed, fuel, or renewable materials, in line with the CFT methodology, the functional unit (FU) is defined as follows:

FU = 1 kilogram of crop produced in a growing season within a crop-country combination

3 Method

The performance measurement approach needed to report on Bayer's commitment to reduce on-field carbon emissions follows the processes highlighted below.

1. Inventory data compilation.
2. Determination of on-field GHG emissions and carbon sequestration with the Cool Farm Tool
3. Calculation of BCS Customer GHG emissions

3.1 Description of the GHG Assessment Inventory data

BCS uses primary inventory data from Kynetec's 'FarmTrak™' which tracks global agriculture in 52 countries, surveying and interviewing from amongst their 300,000 statistically representative grower community annually and collecting details of crop inputs used on over 43 million hectares of land each year. Kynetec, a global agricultural market research company survey customers to collect data needed to estimate GHG emissions using the science-based Cool Farm Tool calculator.

The inventory data used for this study as input for the CFT are sub divided into 2:

(a) Kynetec's FarmTrak primary panel data which focuses on all information related to crop protection, fertilizer and seeds, but lacks data on other farm input domains. These data are collected on an annual basis by interviewing farmers in the relevant markets. FarmTrak™ provides essential information for calculating carbon footprints but lacks data on other input domains.

(b) Kynetec's FarmTrak supplementary sustainability data used for other necessary input information such as soil characteristics, machinery, cultivation techniques etc. This is required to fill the information gaps from the collected Kynetec primary data to enable sustainability-related analyses. Calculating carbon emissions, in collaboration with CFT, is an example of such an analysis.




Figure 2 - Workflow for the calculation of farm level carbon footprints

3.1.1 Sampling approach and processing

One of the aims of FarmTrak™ is to quantify input markets. A representative sampling design that accurately reflects the population is crucial. Kynetec sampling is based on official, statistical data for each crop across regions and is representative of all focus crops on the level of a particular region. The FarmTrak™ samples are built country-by-country while respecting local conditions.

A stratified sampling approach is used when selecting the FarmTrak™ panel respondents. The three elements considered are (1) crop grown (2) location where the crop is grown and (3) size of the farm on which the crop is grown. Consequently, the entire population is split into subgroups considering these criteria. Size of each subgroup is determined by their relative importance in the market. Within each of those subgroups Kynetec applies a random sampling approach, i.e. each respondent belonging to one of these subgroups has the same a priori chance of being interviewed. Quota per subgroup is used and monitored to ensure a representative view of the market. An additional set of criteria are considered when actually selecting the respondents, to ensure Kynetec is interviewing the relevant person. For example, the surveyed respondent must be the farm manager or the person in charge of field level decisions (such as choice of fertilizer, seed, or CP product).




Figure 3 - Qualifying criteria to be met by farmers for selection as part of survey respondents.

FarmTrak™ and the sustainability data collection rely on the same sampling approach, with the only difference being the number of interviews conducted. Usually, no less than one third of the initial FarmTrak™ panel are re-interviewed. In collecting these data, multiple data collection methodologies are deployed such as face-to-face (F2F) interviews, telephone interviews and online surveys. Kynetec achieves a high rate of panel retention thereby ensuring a year-to-year data collection. The consistency of sample

over the years is between 60-90%. However, it is each time a statistically representative sample of randomly selected farmers by Kynetec, a third party independent market research which BCS cannot influence.




Figure 4 - A stratified sampling plan for data collection on Maize (same as corn) cultivation in Belgium.

3.1.2 Data quality check by Kynetec

Several data quality control measures are implemented during and after data collection. First of all, the interviewers attend a professional training course related to research best practices and are given comprehensive instructions on the research procedures. All interviewers are initially accompanied and test-checked for their knowledge and competence.

The following methods are also adopted:

1. Collected data are checked for accuracy and consistency, including (telephone) back-checks.
2. Constant monitoring by fieldwork supervisors checking that all questions are asked correctly, proper responses are recorded, and that interviewers don't need further coaching/training.
3. Tablet and online questionnaires are equipped with proper logic so that farmers only answer questions relevant to them.
4. Based on knowledge and experience in data collection, Kynetec knows the acceptable ranges at product/application level with data collected therefore checks the data against extreme ranges to remove outliers.
5. Identified problem questionnaires are thoroughly reviewed by analysts and are subject to further telephone checks. This survey integrity stage is truly one of the most critical phases of producing this study.
6. Farmers are asked to report all behaviors and decisions. To achieve this, the respondent's anonymity is guaranteed. As a result, the panel data reflects the market realities of some off-label usage that would not otherwise be known.

3.1.3 Inventory data compilation for the GHG assessment

The inventory data compiled for this report are based on such a combination of FarmTrak™ base panel data and supplementary interviews undertaken with ~6,390 panelists in 10 countries for the calculation of carbon emissions. The Table 4 below illustrates the different crop-country combination in which data collection at farm level was focused on for BCS' GHG emission assessment. As a result of the phases in data collection by the data provider, Kynetec, the CCCs are divided into Tier 1 and Tier 2.

Table 4 - Kynetec data collection phases and total sample size for the crop-country-combination's

Tier	CCCs	Harvest year	Total sample size (No of farm surveyed)
1	India-rice, paddy	2020	1000
1	Italy-corn	2020	317
1	Spain-corn	2020	254
1	Argentina-corn	2021	278
1	Australia-cotton	2021	50
1	Brazil-corn	2021	971
1	Brazil-soybeans	2021	926
1	USA-corn	2021	1006
1	USA-soybeans	2021	919
1	USA-cotton	2021	264
2	Canada-rapeseed	2021	324
2	Canada-wheat	2021	268
2	France-wheat	2021	760
2	Mexico-corn	2021	319
2	USA-wheat, spring	2021	269
2	USA-wheat, winter	2021	491
2	Argentina-soybeans	2022	367
2	Australia-wheat	2022	622

BCS argues that such a large data set of farm-level primary data is sufficient for calculating and reporting of crop carbon footprints. For example, Clavreul et al (2017) stated that a minimum of 30 farms is needed for the region of concern for several years' worth of data, particularly as climate change effects become more prevalent and extreme events such as drought or torrential rain becomes norm rather than exception.

3.1.4 CCCs production quantity and BCS market share

The individual CCC production quantity were derived from the Food and Agriculture Organization of the United Nations Statistics (FAO Stats) (<https://www.fao.org/faostat/en/#data/QCL>, accessed on Sept. 27, 2021). This assessment uses the average crop production in the recent five years (2015-2019) for the 18 CCCs. The Table 5 below summarizes the UNFAO reported average crop production in 2015-2019 for the 18 CCCs.

The BCS market share data for each of the 18 CCCs (Table 5) were extracted from the OPTIMAS in 2019. OPTIMAS is an internal BCS market planning tool that provides short- and long-term market planning and market values to serve top management regarding strategic business planning and reporting. The OPTIMAS in 2019 represents the internal market view with market share assumptions that were based in 2019.

The production quantity and the BCS market share are used in deriving the weighting factor (further explanation in section 3.3)

Table 5 - Summary of UNFAO reported average crop production for 2015-2019 and BCS market share for the CCCs.

CCCs	Average crop production (tonnes)	BCS market share
Argentina-corn	44,681,904.0	-. [*]
Argentina-soybeans	53,653,851.6	-. [*]
Australia-wheat	23,274,902.6	-. [*]
Australia-cotton	674,335.5	-. [*]
Brazil-corn	86,177,438.8	-. [*]

Brazil-soybeans	108,154,739.8	-, -*
Canada-rapeseed	19,659,040.0	-, -*
Canada-wheat	30,942,707.0	-, -*
France-wheat	37,354,670.8	-, -*
India-rice, paddy	168,220,346.0	-, -*
Italy-corn	6,488,032.8	-, -*
Mexico-corn	27,021,010.2	-, -*
Spain-corn	4,087,309.8	-, -*
USA-corn	368,030,854.0	-, -*
USA-soybeans	112,251,616.0	-, -*
USA-wheat, spring	10,849,670.5	-, -*
USA-wheat, winter	41,852,356.6	-, -*
USA-cotton	3,775,907.5	-, -*

* The Bayer Crop Science Division market share data extracted from our internal market planning tool were shared with the panel of experts under a non-disclosure agreement.

3.1.5 Definition of BCS customer base used for the on-field GHG assessment

For the on-field GHG assessment, BCS uses compiled inventory data for all 18 CCCs until the end of 2022. The GHG performance are measured and aggregated on the CCC level (for CCC-specific baseline values), and a consolidated global GHG performance across all CCCs selected (for a global aggregated baseline value) is calculated.

The BCS GHG target is measured as a 30% reduction of on-field GHG emissions per kilogram of crop produced by BCS's farming customers in 2030 for the most emitting crop systems in the regions where BCS operates. Therefore, the focus of the BCS on-field GHG commitment is on the GHG emissions and carbon sequestration of BCS's farming customer base (i.e., specific field gate-to-gate emissions per kg of crop weight) for any BCS's farming customer in a particular CCC.

Because farmers in the FarmTrak™ panel data might use solutions from different competitors simultaneously, BCS's farming customers will be identified and distinguished in FarmTrak™ following the below mentioned reasoning. The farms will be identified relying on "share of wallet" calculations, comparing it with BCS's market share in a CCC.

Farmers are considered to be BCS's customers based on the following principles:

1. BCS's share of wallet of a particular farm at least equals BCS's market share for the relevant country/crop combination (see equation 1 and 2 below) and / or
2. They use BCS's seed variety and / or
3. They use BCS's 'Climate Field View⁶' or any other digital platform from Bayer and / or
4. They are being incentivized by BCS for adoption of climate-smart practices by participating in our carbon business models

$$\text{Market share (in a CCC)} = \frac{\text{Hectares treated with Bayer products in entire market}}{\text{Total hectares treated with crop protection in entire market}} \quad (1)$$

Market share per country will be calculated considering hectares treated with Bayer's products relative to total hectares treated with crop protection in that market. Market share will be calculated considering all product lines. Hectares treated refers to "Super Developed Area" and takes multiple applications on

⁶ Climate Field View is BCS's digital farming software platform that helps farmer to monitor and make agronomic decisions on their fields for yield optimization and profit maximization.

same field into account. For example: if a field of 10 hectares is treated twice, BCS considers hectares treated/super developed area to be 20 hectares.

$$\text{Share of wallet (on a farm)} = \frac{\text{Hectares treated with Bayer products on the farm}}{\text{Total hectares treated with crop protection on the farm}} \quad (2)$$

In the current calculations, share of wallet states how much respondents spend/use on Bayer's products exclusively. Share of wallet can be calculated considering hectares treated with Bayer products relative to total hectares treated on the same farm.

Share of wallet allows to evaluate how Bayer is performing against competitors and allows to benchmark against Bayer's market share of a particular country-crop combination. All farms will be identified as Bayer customers if Bayer's share of wallet of a particular farm at least equals Bayer's market share for the relevant country-crop combination. Farms will be identified as non-customers, if Bayer's share of wallet of a particular farm is smaller than Bayer's market share for the relevant country-crop combination.

Table 6 - Sample size and number of customers according to definition of BCS's customer base.

CCC	Total sample size (No of farm surveyed)	No. of BCS customers
Argentina-corn	278	124
Argentina-soybeans	367	172
Australia-wheat	622	217
Australia-cotton	50	32
Brazil-corn	971	611
Brazil-soybeans	926	602
Canada-rapeseed	324	236
Canada-wheat	268	158
France-wheat	760	298
India-rice, paddy	1000	258
Italy-corn	317	185
Mexico-corn	319	232
Spain-corn	254	168
USA-corn	1006	600
USA-soybeans	919	507
USA-wheat, spring	269	134
USA-wheat, winter	491	195
USA-cotton	264	129

3.2 Determination of on-field GHG emissions and carbon sequestration with the Cool Farm Tool

3.2.1 Cool Farm Tool model description

The Cool Farm Tool (CFT) is an online greenhouse gas (GHG) calculator that quantifies the carbon footprint of crops in kgCO₂ equivalents (kgCO₂e) over a 100 year time horizon. The tool has a specific farm-scale, decision-support focus making it possible to identify emissions on the field. It further provides farmers with

the opportunity to evaluate different management options that will lead to positive impact on the total emissions from the farm. Contrary to other farm GHG emission calculators, it includes a calculation of soil carbon sequestration which is an important aspect of agriculture GHG accounting in terms of adaptation and mitigation benefits. As a result of its use of readily available farm data, there is considerable scope for its use in global surveys to inform on current practices and potential for mitigation (Hillier, et al., 2011).

The CFT was originally developed by Unilever and researchers at the University of Aberdeen to help growers measure and understand on-farm GHG emissions. The use of the tool is designed to be simple, but scientifically robust in accounting for farm GHG emissions. It has been tested and adopted by many multinational companies which are using it to work with farmers to measure, manage and reduce GHG emissions arising from crop production towards contributing to the mitigation of climate change. More information about CFT can be found at <http://www.coolfarmtool.org>.

The CFT was selected for this assessment because of its ease of use, widespread adoption, global applicability, decision-support focus and its readily availability of farm data for the intended purpose which is to calculate GHG emissions. The CFT is being used by diverse array of stakeholders which includes food retailers, manufacturers, input suppliers, NGOs, universities and consultancies. A list of CFT partner members can be found at <https://coolfarmtool.org/cool-farm-alliance/members/>.

The methodology used in the CFT, calculates GHG emissions and removals associated with the production of an agricultural product. A carbon footprint is reported for the three major sources of on-farm emissions associated with the production of agricultural products, namely, carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O). For crops, the CFT incorporates IPCC Tier⁷1, Tier 2 and when it comes to N₂O emissions and soil carbon sequestration, a simplified Tier 3 multi-factorial empirical model based on Bouwman et al. (2002), which is widely acknowledged is used. The Cool Farm Tool is moving towards Tier 3 when possible. Currently, the CFT is based on the IPCC 2019 refined guidelines and uses the Global Warming Potentials from the IPCC Assessment Report 6 (Cool Farm Alliance, 2022). Detailed information on the data needed to calculate GHG emissions from crops is summarized in the CFT data input guide (Cool Farm Alliance, 2021). Please refer to the [CFT data input guide](#), the [CFT FAQ](#), and Hillier et al., (2011) for a detailed technical description of the CFT methodology.

The CFT has several input sections which are listed below. Each section requires provision of information related to the crop being assessed. The carbon footprints are calculated for one selected growing area/parcel/field per farm, assuming similar soil characteristics and input/management practices on that same area/parcel/field. For each crop and growing area, a full annual production cycle is considered. The scope of the current project is to consider emissions before the crop leaves the farm (i.e., everything on-field before 'farmgate').

3.2.2 Cool Farm Tool input data

The CFT is structured according to the following sections:

- (0) Farm settings
- (1) Crop
- (2) Soil
- (3) Inputs
- (4) Fuel & Energy
- (5) Irrigation
- (6) Carbon
- (7) Transport (excluded)

When using the CFT for emission calculation, some input parameters have been predefined in the model while some are to be defined by the user. In the next sub-chapters, we will go into the details of the input

⁷ A tier represents a level of methodological complexity used in GHG calculation. There are three tiers namely Tier 1, Tier 2, Tier 3. Tier 1 is the basic method, Tier 2 represents the intermediate while Tier 3 is the most complex in terms of the methodology.

parameters used in the calculation of GHG emissions. The BCS inputs used in the next sections are based on Kynetec data (see section 3.1 for details).

0. Farm settings

This is the base section where details about the farm location and the climate condition are defined. The input parameters are described in Table 7 below.

Table 7 - Cool Farm Tool Input parameters on farm settings

Variable	Options / Unit	Description
Country	-	Country where the farm is located.
Annual average temperature	°C	This information is not collected during the interview with farmers, instead Kynetec relies on external sources (such as the National Oceanic and Atmospheric Administration - NOAA)
Climate	- Temperate - Tropical	Climate zones are defined following the logic of Bouwman et al. (2002) who categorize all Global Ecological Zones (FAO, 2010) as either temperate or tropical: - Tropical: tropical and subtropical - Temperate: temperate and boreal

1. Crop

This section is divided into three input sections which are crop details, crop residue management and co products.

1.1 Crop details

Information here includes the type of crop, area for crop growing and the crop yield (see Table 8 below for details on the Input data required for crop details). The CFT has an additional emission calculation for rice when cultivated as paddy rice. This is because paddy rice plays a significant role in the overall emission from agriculture. The CFT accounts for the emission from paddy production using the IPCC approach based on Yan et al. (2005). The emission factor from this approach considers water regime during cultivation, water regime in the pre-season and organic amendments. The current version of CFT only covers seed emissions for potato, but not for other field crops. The model considers emissions from seeds to be quite low, compared to the other sources of emission.

Table 8 - Cool Farm Tool Input parameters on crop details

Variable	Options / Unit	Description
Crop name	-	Name of the crop.
Harvest year	-	Calendar year during which the crop was harvested.
Crop area	Hectare	Size of the parcel, including buffer zones.
Harvested amount	Metric ton	Total harvested crop from the crop area for the relevant harvest year before on-farm processing (eg drying, grading, sorting) of crops i.e. Fresh matter
Farm-gate ready amount	Metric ton	Total marketable yield from the crop area for the relevant harvest year after on-farm processing.
Assessment name	-	A reference name for the identification of the assessment.

1.2 Crop residue management

Crop residue refers to the plant matter from crop production that is not used as a sellable product. Often, harvest does not cover the full biomass of a crop and thus crop biomass remains as residue both above and below ground. Examples of residue from crop production typically include leaf lamina, leaf mid-rib, pseudostem sheath, fruit peelings etc.

For the calculation of emissions from residue in the CFT, the amount of residues generated per year and the way residues are managed are required as input data. If residues are used to create compost, the tool will calculate the possible emissions associated with this compost production process. However, if compost is then used on crops, an emission factor of zero is associated with the compost since it is already accounted for in the residue section. When residues are used as compost, the emission increases depending on the technology (forced aeration or non-forced aeration) used during composting. Non-forced aeration accounts for more emission compared to forced aeration. Detailed description of the required input can be found in Table 9 below

Table 9 - Cool Farm Tool Input parameters on crop residue management

Variable	Options / Unit	Description
Residue amount	Ton/ha	The default residue amount estimated by the CFT for various crops is used in this assessment.
Residue management	-	<p>The CFT provides the following pre-defined options for selection.</p> <ul style="list-style-type: none"> - Removed from field for use or for sale. - Used for composting: Forced aeration or non-forced aeration compost. - Left untreated in heaps. - Burnt on the field. - Distributed on the field, incorporated or mulched. <p>The above options are selected for the assessment based on the responses from the farmers on how they manage crop residues.</p>

1.3 Co-products

This section of the CFT allows allocating the total crop emissions between main product (e.g. wheat) and co-product (e.g. straw). However, BCS excludes co-products because it does not allocate a proportion of emissions of the main crop to one or more co-products. The estimated GHG emissions from co-products are associated with main product. Therefore, this assessment uses the default by allocating all emissions to a single main product.

2. Soil

This section is where the soil characteristics of the field being assessed are specified. In defining the soil characteristics, the CFT considers input from the soil texture, soil organic matter, soil moisture, soil drainage and soil pH. The pre-defined chosen input range for soil organic matter is used in determining the soil organic carbon. Detailed description of the required input can be found in Table 10 below.

Table 10 - Cool Farm Tool Input parameters on soil characteristics

Variable	Options / Unit	Description
Soil texture	<ul style="list-style-type: none"> - Fine - Medium - Coarse 	<p>Soil texture is based on soil type, as stated by the grower and grouped accordingly:</p> <ul style="list-style-type: none"> - Fine: sandy clay, clay, silty clay - Medium: sandy clay loam, clay loam, silty clay loam - Coarse: sand, loamy sand, sandy loam, loam, silt loam, silt

Soil organic matter	<ul style="list-style-type: none"> - SOM \leq 1.72% - 1.72% < SOM \leq 5.16% - 5.16% < SOM \leq 10.32% - SOM > 10.32% 	The soil organic matter is expressed as percentage. As stated by the grower, selection is made based on the four categories.
Soil moisture	<ul style="list-style-type: none"> - Moist - Dry 	As stated by the grower. Moist soils are those without any water constraints during the growing season.
Soil drainage	<ul style="list-style-type: none"> - Good - Poor 	As stated by the grower. Soils which are often saturated or show surface water were classified by the grower as 'Poor', other soils are classified as 'Good'.
Soil pH	<ul style="list-style-type: none"> - pH \leq 5.5 - 5.5 < pH \leq 7.3 - 7.3 < pH \leq 8.5 - pH > 8.5 	As stated by the grower, selection is made based on the four categories.

3. Inputs

This section is divided into two input sections which are fertilizer inputs and crop protection inputs. These inputs have influence on the GHG emission of the farm. The emissions resulting from the fuel used in applying of these inputs are entered in the 'Fuel & Energy' section. Detailed description of how emissions from fertilizer and crop protection inputs are calculated are described below:

3.1 Fertilizers

In the case of fertilizers, the CFT accounts for two types of emission pathways: emissions released during fertilizer manufacturing and emissions from the application of fertilizer on the field. Since emissions from fertilizer manufacturing are considered out of scope for the BCS on-field GHG commitment, only emissions from the application of fertilizer on the field are covered: These emissions are mainly triggered by bio-chemical process related to the addition of nitrogen fertilizers and limestone. Although emissions from soils may happen without the use of fertilizer, fertilizer application is one of the major sources of N₂O emissions. From the input of the type of fertilizer used on the field, the CFT tool defines the N:P: K ratio of the fertilizer. For nitrous oxide (N₂O) and nitric oxide (NO) emissions resulting from nitrification and denitrification process, the factor values from the multivariate empirical model of Bouwman et al. (2002) were used. NO and NH₃ emissions are converted to N₂O using recommended IPCC factor. Volatilization of NH₃ is also taken into account using the equation from FAO and IFA (IFA and FAO, 2001), and the recommended IPCC conversion factor is used for NH₃ to N₂O. In moist soils, some of the added Nitrogen fertiliser are lost through leaching. Factors from IPCC are used to estimate the amount of Nitrogen that are lost through this pathway and the resulting N₂O emissions. The emission effect from the presence of nitrification inhibitors in fertilizers are modelled using the methodology by Akiyama et al., (2010). The CFT methodology used in accounting for emissions associated with field application of fertilizers takes into account the different types of fertilizers, crop type, soil properties and fertilizer application methods.

Table 11 - Cool Farm Tool Input parameters on fertilizer management

Variable	Options / Unit	Description
Fertilizer type	Pre-defined list of applicable fertilizers	As stated by the grower, the fertilizer used during crop production is selected here from the CFT predefined list.
Application rate	Kg or L per Hectare	The amount of fertilizer used per hectare, as stated by the growers
Fertilizer weights or units	Product or Units of active element	Units of product (kg or liter) is used as default option.
Application method	<ul style="list-style-type: none"> - Broadcast - Incorporate - Apply in solution 	As stated by the grower, a selection is made on how the fertilizer is applied on the field.

	- Fertigation	
Emission inhibitors	- None - Nitrification inhibitor	For each fertilizer applied, the growers mentions if the fertilizer contains an emission inhibitor or not. None is chosen when the applied fertilizer contains no inhibitor.

3.2 Crop protection inputs

The CFT assumes that a part of the emissions from use of CPP occurs during their production. Since this type of embodied emissions take place off- field, they are out-of-scope and not considered in this report. Emissions related to the energy use from applying the crop protection products on the field are accounted for in the direct energy section.

4. Fuel & Energy

This section deals with the estimation of emission resulting from energy consumption in the growing area. Possible energy sources that are considered are electricity and fuels. This includes on site energy use for machinery and irrigation. The consumption of fuel and the use of energy for farm operation adds to the overall emissions from agricultural production. The emission calculation includes both electricity and liquid fuel use. For energy sources which consist of diesel, petrol, bioethanol, biodiesel, electricity (grid, hydroelectricity and wind), the CFT uses emission factors derived from the GHG protocol (2003). The CFT does not assume a zero emissions factor for renewable energy. Emissions for electricity from renewable energy are significantly lower than for electricity from the grid but not accounted as zero due to emissions released during the development of renewable energy technology and construction of plants. In situations when the data of annual amounts of energy sources consumed for certain activities are not available, indirect figures such as number of applications, machinery/vehicle type, fuel type, and size of area treated are used to compute emissions.

In the CFT, this section is divided into three parts: Direct energy use, field operations energy use and wastewater.

4.1 Direct Energy

Energy consumption related to irrigation is accounted for in direct energy use. See Table 12 below for details on the input parameters for this section

Irrigation

Two steps are undertaken to estimate energy consumption from irrigation pumps:

- Desk research⁸ is done to estimate irrigation volumes at relevant subnational level (i.e. state, province etc.), focusing on the most recent statistics available. Different methods of irrigation (i.e. flooding, rain gun, pivot or drip irrigation) are considered. (For example, EASAC, USDA-NASS, ANA Brazil etc.)
- The CFT calculates the energy requirements in kWh for irrigating 1 mm/ha depending on irrigation method and fuel used. These reference values are used to estimate energy consumption.

Table 12 - Cool Farm Tool Input parameters on direct energy use

Variable	Options / Unit	Description
Energy source	Predefined list of different sources of energy	Electricity or diesel is assumed to be the relevant energy sources.

⁸ Alternatively, water irrigation volumes can be estimated using the 'water footprint calculator' from CFT. At the time the data was processed this was not available yet but can be considered for future data processing.

Energy used	- Kwh - liter	Volume of energy used (liter of diesel or KWH electricity)
Category	- Field - Facility	Energy consumption from irrigation is categorized as 'field'.

4.2 Field Operations Energy Use

Energy consumption related to on-field machinery operations is taken into account in the section 'field operations energy use.' The CFT supports estimating fuel use for common agricultural machinery from tillage, sowing, spraying crop protection, fertilizer applications and harvesting. The focus of this section is to determine energy used based on machinery operation on the field. Required inputs are the type of machine (obtainable from a pre-defined list), fuel used and number of field operations. Type and number of field operations are entered following the below mentioned logic.

- Sowing and cultivation practices: As part of the 'sustainability' data collection, growers are asked to mention which one of three cultivation practices they adhere to (1) conventional tillage (2) reduced tillage (3) zero tillage. Building on the logic as described in Khaledian et al. (2014) these cultivation practices result in the below mentioned machinery operations. These are mapped accordingly on the CFT machinery typology.

Table 13 - Cool Farm Tool Input parameters on cultivation practices and field operations

Cultivation practice	Machinery operations
Conventional tillage	Plowing, Harrow, Disc Harrow, Seed Drill
Reduced tillage	Harrow, Disc Harrow, Seed Drill
Zero tillage	No-till Seed Drill

- CPP spraying and fertilizer applications: Number of times the field was visited for applying crop protection products and fertilizers is derived from the FarmTrak™ crop protection data and sustainability data. Both databases provide information on the timing of the different applications. All applications that happen on the same date are aggregated and are assumed to happen during one single pass for fertilizers and crop protection (cf. concept of tank mix for crop protection data).
- Harvesting and residue management: Kynetec assumes that harvesting is mainly done with a combine (e.g. cereals, soybean, corn), or could be done manually in some smallholder markets (e.g. India rice). A special 'Corn combine' is selected for harvesting corn. In case the grower mentioned that the crop residue is taken off field, a pass with a baler for collecting the residue is added.

Table 14 - Cool Farm Tool Input parameters on field operations energy use

Variable	Options / Unit	Description
Machine category	Predefine list of different farm operation.	Selection is made based on different farm operations. Eg Harvesting, tillage, spraying, sowing, fertilization.
Machine	Predefine list of different machines based on the selected farm operation	Selection is made based on different machines used in farm operations. For example, when spraying was selected as machine category, herbicide sprayer was selected here.
Fuel use	- Diesel - Petrol	Diesel is used as a default fuel type for machinery.
Number of operations	-	Number of completed field operations related to the farm operation being assessed during the growing cycle for the crop. Filled based on response from the growers

4.3 Wastewater Emissions

Most crops do not have wastewater emissions and are thus not accounted for in GHG emission calculation. Methane emissions from wastewater arises from the decomposition process of organic material. This is common in coffee where a wet milling process is used to separate the pulp from the bean. The Bayer CCC has no coffee as part of the selected crops, therefore wastewater emission is not relevant for this report.

5. Irrigation

In the irrigation section, a repeated computation of irrigation energy was not carried out as energy for operating irrigation pumps is already captured in the 'Direct energy' section.

6. Carbon

This section describes the emission resulting from changes in management practices that alters the carbon stocks i.e. carbon stored by or released from the soil and above ground biomass of the growing area. Changes in carbon stocks can occur from alterations in land use, soil management, and biomass. They can affect net carbon capture or release, thereby impacting emissions. Land use change (e.g. deforestation) is not considered in this report (see section 5 on limitation for more information). Soil management practices considered are tillage and cover crops. Management changes can either increase or decrease the carbon in the soil and will continue doing so until a new equilibrium is reached. The CFT only considers changes in farm management practices that have occurred within the last 20 years because this time frame is assumed by IPCC and other GHG accounting standards as the period that soil carbon stocks need in order to reach a new equilibrium. Any management change that has happened before is assumed to be no longer relevant.

In the CFT, determination of the carbon stocks in the top 30 cm of the soil are based on the user soil characteristics input and are determined mathematically using bulk density and carbon density. The carbon density describes the carbon available in the top 30 cm of 1 ha of soil based on an assumed bulk density of 1 g/cm³ and 1% soil organic matter equals 1.72% of soil organic carbon. The IPCC Tier 1 method is used for the estimation of soil carbon stock changes using coefficients from (Ogle S.M., 2005) for carbon stock changes related to change in management practice for a period of 20 years. The resultant amount of a change in soil carbon is dependent on climate (Hillier, et al., 2011). The changes in carbon were converted to an annualized CO₂ emission (can either be positive or negative) when land management changes in relation to carbon input practice and tillage practice. The carbon input practice is classified into low, medium and high. Low refers to minimal residue return as a result of residue removal, medium category accounts for annual cropping with cereals where residues are returned to the field while high is in addition to medium with higher inputs due to production of high residue yielding crops, cover crops, improved vegetated fallows and frequent use of perennial grasses in annual crop rotations. The tillage classes (conventional, reduced, or no till) are defined following IPCC classification. The changes in soil carbon stock as a result of manure and compost addition are derived from Smith et al. (1997).

Table 15 - Cool Farm Tool Input parameters on tillage and cover crops management

Variable	Options / Unit	Description
Changed from ...		Based on the information Kynetec gets from the growers, these types of management change are considered: <ul style="list-style-type: none">- Tillage: Comparison of how the field was tilled (conventionally, reduced or not)- Cover crops: Checking if a cover crop is grown
Number of years ago	-	Number of years ago the situation changed.
Percentage of field	%	For changes related to tillage and cover crops, it is assumed that it happened on the entire field.

3.3 Calculation of BCS Customers GHG emission

In the following section, we describe in detail the methodology in the calculation of BCS Customer GHG emission for the baseline year. This section further includes the formulae that will be used for future tracking of the performance.

3.3.1 Calculation of specific GHG baseline for particular CCC's

The specific GHG emission (kg CO₂e per kg crop) is the **normalized gate-to-gate GHG emissions calculated for an individual CCC**. For a baseline and a specific base year, the specific GHG emission is calculated as shown in equation 3 below:

$$GHG_{BL,spec}^{CCC} = \frac{\sum_{i=1}^k GHG_{i,BL}^{CCC}}{\sum_{i=1}^k W_{i,BL}^{CCC}} \quad (KgCO_2/KgCrop)$$

3

For k farmers assessed in a base year for a particular CCC:

- $GHG_{BL,spec}^{CCC}$ = Specific GHG emissions for a particular CCC in the base year
- $GHG_{i,BL}^{CCC}$ = Absolute GHG emissions of a farmer i for a particular CCC in the base year
- $W_{i,BL}^{CCC}$ = Crop weight (Kg) of a farmer i for a particular CCC in the base year

To track GHG reductions over time until 2030, the **baseline GHG performance in a particular CCC will be compared with the GHG performance in a future target year (t)** (e.g. next performance tracking year 2024) based on the following formula:

$$GHG_{t,spec}^{CCC} = \frac{\sum_{i=1}^n GHG_{i,t}^{CCC}}{\sum_{i=1}^n W_{i,t}^{CCC}} \quad (KgCO_2e/kgCrop)$$

4

For n farmers assessed in a target year t for a particular CCC:

- $GHG_{t,spec}^{CCC}$ = Specific GHG emissions for a particular CCC in a year t
- $GHG_{i,t}^{CCC}$ = Absolute GHG emissions of a farmer i for a particular CCC in a year t
- $W_{i,t}^{CCC}$ = crop weight (kg) of a farmer i for a particular CCC in a year t

As the absolute emissions and crop weight values are separately summed up, specific BCS emissions are weighted according to different crop weights and, indirectly, field sizes.

Finally, a specific GHG emission reduction is calculated as:

$$R_t^{CCC} = \left[1 - \frac{GHG_{t,spec}^{CCC}}{GHG_{BL,spec}^{CCC}} \right] \times 100 \quad (\%)$$

5

- R_t^{CCC} = Specific GHG emission reduction for a particular CCC in a year t as compared with the base year

Additionally, target achievement for this particular CCC can be calculated as:

$$TA_t^{CCC} = \left[\frac{R_t^{CCC}}{T^{CCC}} \right] \times 100\% \quad (\%)$$

6

- TA_t^{CCC} = Target achievement for a particular CCC in a year t
- T^{CCC} = Target set for a particular CCC

3.3.2 Setting an aggregated baseline for GHG emission reduction across CCCs

To calculate GHG emissions across CCCs for a baseline and a particular year (for an aggregated global baseline value), the individual baseline results which are specific for each CCC (as described above) need to be aggregated. For this aggregation, the specific baseline GHG emissions for a particular CCC in the base year ($GHG_{BL,spec}^{CCC}$) are weighted with a weighting factor (Wf_{CCC}) which is also specific for each CCC.

$$GHG_{BL,agg} = \sum_{CCC} GHG_{BL,spec}^{CCC} \times Wf_{CCC} \quad (KgCO2e/kgcrop)$$

7

- $GHG_{BL,agg}$ = Aggregate GHG baseline emissions across CCCs (weighted to represent BCS market)
- $GHG_{BL,spec}^{CCC}$ = Specific GHG emissions for a particular CCC in the base year
- Wf_{CCC} = Weighting factor for a particular CCC in the base year

The weighting factors (Wf_{CCC}) are determined by the total production volume of a particular crop in a particular market multiplied by BCS market share and by the specific GHG footprint of BCS customers in this CCC (baseline). The combination of the production volume, the BCS market share and the specific GHG emissions is referred to as the Total GHG emission (**kg CO₂e**). To avoid complexity, these weights are determined once during baselining and then kept fixed⁹ (for the future).

$$Wf_{CCC} = \frac{P_{CCC} \times M_{CCC} \times GHG_{BL,spec}^{CCC}}{\sum_{CCC} P_{CCC} \times M_{CCC} \times GHG_{BL,spec}^{CCC}} = \frac{Total\ GHG}{sum\ of\ Total\ GHG\ across\ CCC's} \quad (unitless)$$

8

- Wf_{CCC} = Weight of a particular CCC in the portfolio (determined during baselining and fixed) (dimensionless)
- P_{CCC} = Production volume of a particular crop in a particular market (FAO database) (tonne)
- M_{CCC} = BCS' market share in a particular market (%)
- $GHG_{BL,spec}^{CCC}$ = Specific GHG emissions for a particular CCC in the base year (kgCO₂e / kgCrop)

Also, for future target years, the specific GHG emissions for a particular CCC in a year t ($GHG_{t,spec}^{CCC}$) will be weighted with the fixed weighting factor (Wf_{CCC}).

$$GHG_{t,agg} = \sum_{CCC} GHG_{t,spec}^{CCC} \times Wf_{CCC} \quad (kgCO2e/kgcrop)$$

9

- $GHG_{t,agg}$ = Aggregate GHG emissions across CCCs in a year t (weighted to represent BCS market)
- Wf_{CCC} = Weight of a particular CCC in the portfolio (determined during baselining and fixed)
- $GHG_{t,spec}^{CCC}$ = Specific GHG emissions for a particular CCC in a year t

Finally, a specific (i.e. relative) GHG emission reduction is calculated across CCCs as:

$$R_t = \left[1 - \frac{GHG_{t,agg}}{GHG_{BL,agg}} \right] \times 100 \quad (\%)$$

10

⁹ Note: Base year is CCC-specific.

- R_t = Specific (i.e. relative) GHG emission reduction across CCCs in a year t as compared with the baseline

Additionally, target achievement across CCCs can be calculated as:

$$TA_t = \left[\frac{R_t}{30\%} \right] \times 100 \quad (\%)$$

- TA_t = Target achievement across CCCs in a year t at the overall target of 30%

11

The performance will be tracked by frequently collecting data, calculating the GHG performance in future years based on the same methodology as described in the above sections, and then comparing the future performance with the baseline performance.

Following guidance from the Greenhouse Gas Protocol Corporate Accounting and Reporting Standard (WRI, World Resources Institute, 2004), for consistent tracking of emissions over time, the base year emissions may need to be retroactively recalculated/restated as BCS undergo significant structural changes such as:

- Inclusion or exclusion of crop-country combinations.
- Investments or divestments.
- Change of boundaries.
- Changes in calculation methodology or improvements in the accuracy of emission factors or activity data that result in a significant impact on the base year emissions data.
- Discovery of significant errors, or a number of cumulative errors, that are collectively significant.

Consequently, BCS shall develop a base year emissions recalculation policy, and clearly articulate the basis and context for any recalculations. If applicable, the policy shall state any 'significance threshold' for deciding on historic emissions recalculation. A significance threshold is a quantitative criterion used to define any significant change to the data, inventory boundary, methods, or any other relevant factors (WRI, World Resources Institute, 2004).

It is the responsibility of BCS to determine the 'significance threshold' that triggers base year emissions recalculation and to disclose it. Based on recommendations of the California Climate Action Registry, the change threshold is set to 10 percent of the overall specific base year emissions, determined across CCCs from the time the base year is established.

In sum, if BCS realizes in the future that significant structural changes as described above happen, BCS will re-check the baseline performance value. If the re-checked baseline performance value differs by 10% from the currently calculated baseline value, BCS will restate the baseline and re-evaluate the further implications for the progress tracking towards the 30% reduction commitment.

4 Interpretation

4.1 Results and setting of GHG emission tracking baseline.

Towards achieving the 30% reduction in the on-field GHG emissions per kg of crop in our customer base, first will be to determine the baseline upon which our progress will be tracked. We calculated the specific GHG emission based on the gate-to-gate GHG emissions using the CFT and the crop weight from the surveyed customer farms in the 18 CCC's. The specific GHG emission is the same as GHG footprint, with a unit of kg CO₂e per kg crop weight. We then calculated the total GHG emissions based on the specific GHG emissions, United Nations Food and Agriculture Organization (UN FAO) reported 5-year average crop production in 2015-2019 (UNFAO 2021) and BCS market share for each of the 18 key markets. The

baseline calculation of the GHG emissions was carried out by Kynetec using the CFT version 1.0 with results described in the following sections

4.1.1 Specific GHG emissions for the baseline

The baseline specific GHG emissions varied across the 18 CCC's ranging from 0.07 kg CO₂e per kg crop in Italy-Corn to 0.97 kg CO₂e per kg crop in India-Rice (Table 16, Figure 5 below). Corn (0.07- 0.13 kg CO₂e per kg crop in Italy, USA, Spain, Argentina) had smallest specific emissions compared to other crops, expect for Brazil-corn (0.18 kg CO₂e per kg crop) and Mexico-corn (0.42 kg CO₂e per kg crop). USA-soybean had the specific emission of 0.14 kg CO₂e per kg crop and France-wheat had the specific emission of 0.19 kg CO₂e per kg crop), then USA-wheat winter had the specific emission of (0.25 kg CO₂e per kg crop). Brazil and Argentina soybean (0.26, 0.38 kg CO₂e per kg crop respectively), US-spring and Canada wheat (0.29, 0.32 kg CO₂e per kg crop respectively) and Canada rapeseed (0.45 kg CO₂e per kg crop) had moderate specific emissions. While cotton had higher specific emissions (0.56 kg CO₂e per kg crop in USA and Australia) followed by Australia-wheat with a specific emission of 0.61 kg CO₂e per kg crop mainly contributed by the smallest total crop weight. Rice in India had the largest specific emission (0.97 kg CO₂e per kg crop) of all the 18 CCCs.

Table 16 - Specific GHG emission (kg CO₂e per kg crop) from surveyed BCS customer farms in the baseline

CCCs	CFT- modeled GHG emission (kg CO ₂ e)	Crop weight of surveyed customer (Tonne)	Specific GHG emission (kg CO ₂ e per kg crop)
Argentina-corn	148839197	1109361.9	0.1342
Australia-cotton	5236418.46	9408.22	0.5566
Brazil-corn	135548634.1	774304.77	0.1751
Brazil-soybean	150393044.6	584936.18	0.2571
India-rice	3526317.05	3634.71	0.9702
Italy-corn	4612713.74	64558.71	0.0715
Spain-corn	10641954.47	95512.85	0.1114
USA-corn	272990301.3	2909794.63	0.0938
USA-cotton	49354335.23	88910.75	0.5551
USA-soybean	94663958.61	683436.8	0.1385
Argentina-soybean	130193457.4	344495.7	0.3779
Australia-wheat	4775618.29	7877.66	0.6062
Canada-rapeseed	33124891.18	75533.29	0.4385
Canada-wheat	19814089.57	61022.7	0.3247
France-wheat	6221656.53	32100.5	0.1938
Mexico-corn	26203826.23	62709.07	0.4179
USA-wheat, Spring	13184910.7	45565.18	0.2894
USA-wheat, Winter	13481567.08	54669.41	0.2466




Figure 5 - Specific GHG emission in the baseline based on BCS'S customer farms

The Emission sources contributing to GHG emissions varied greatly. Fertilizer decomposition was a major contributor to GHG emissions across the 18 CCC's, except for India rice. In the India rice market, methane emissions from the paddy play a major role in GHG emissions. (see Table 19 in the appendix for the numbers and see below Figure 6 for the graph on the different emission sources).




Figure 6 - Specific GHG emission in the baseline based on BCS'S customer farms grouped by emission sources

4.1.2 Total GHG emissions across the CCC for the baseline

The total GHG emissions in the baseline varies across the 18 markets (Table 17, Figure 7). This is due to market share, specific GHG emission and the production amount. Spain-corn had the smallest total emission followed by Italy-corn and India-rice had the largest total emission, mainly contributed by the highest specific GHG emission. USA-corn has the second largest total emission, mainly contributed by the largest crop production.

Table 17 - Total GHG emissions (kg CO₂e) in the baseline year and different factors used in calculating Bayer's total GHG emissions in the 18 CCCs.

CCCs	Average Crop Production (tonne)	Bayer's Market Share	Specific GHG Emission (kg CO ₂ e per kg crop)	Total GHG Emission (Kg CO ₂ e)
Argentina-corn	44681904	-.*	0.1342	-.*
Australia-cotton	674335.5	-.*	0.5566	-.*
Brazil-corn	86177438.8	-.*	0.1751	-.*
Brazil-soybean	108154739.8	-.*	0.2571	-.*
India-rice	168220346	-.*	0.9702	-.*
Italy-corn	6488032.8	-.*	0.0715	-.*
Spain-corn	4087309.8	-.*	0.1114	-.*
USA-corn	368030854	-.*	0.0938	-.*
USA-cotton	3775907.5	-.*	0.5551	-.*
USA-soybean	112251616	-.*	0.1385	-.*
Argentina-soybean	53653851.6	-.*	0.3779	-.*
Australia-wheat	23274902.6	-.*	0.6062	-.*
Canada-rapeseed	19659040	-.*	0.4385	-.*
Canada-wheat	30942707	-.*	0.3247	-.*
France-wheat	37354670.8	-.*	0.1938	-.*
Mexico-corn	27021010.2	-.*	0.4179	-.*
USA-wheat, Spring	10849670.5	-.*	0.2894	-.*
USA-wheat, Winter	41852356.6	-.*	0.2466	-.*




Figure 7 - Total GHG emission based on BCS's customer farms for the 18 CCC's

4.1.3 Aggregated GHG emission for the baseline year

To determine the overall baseline, the individual CCC baselines are weighted. The weighting factors for each CCC is the ratio of total emissions in each market to total emissions across all the markets. They vary greatly across key market (Table 18, Figure 8). Based on weighting factors, the weighted specific GHG emission across 18 key markets in the baseline is **0.443 kg CO₂e per kg crop**.

Table 18 - The specific GHG emission, total GHG emission and weighting factors for the 18 CCCs

CCCs	Specific GHG Emission (kg CO2e per kg crop)	Total GHG Emission (kg CO2e)	Weighting Factor
Argentina-corn	0.1342	-.-*	-.-*
Australia-cotton	0.5566	-.-*	-.-*
Brazil-corn	0.1751	-.-*	-.-*
Brazil-soybean	0.2571	-.-*	-.-*
India-rice	0.9702	-.-*	-.-*
Italy-corn	0.0715	-.-*	-.-*
Spain-corn	0.1114	-.-*	-.-*
USA-corn	0.0938	-.-*	-.-*
USA-cotton	0.5551	-.-*	-.-*
USA-soybean	0.1385	-.-*	-.-*
Argentina-soybean	0.3779	-.-*	-.-*
Australia-wheat	0.6062	-.-*	-.-*
Canada-rapeseed	0.4385	-.-*	-.-*
Canada-wheat	0.3247	-.-*	-.-*
France-wheat	0.1938	-.-*	-.-*
Mexico-corn	0.4179	-.-*	-.-*
USA-wheat, Spring	0.2894	-.-*	-.-*
USA-wheat, Winter	0.2466	-.-*	-.-*





Figure 8 - Individual weighting factors for the 18 CCCs

4.1.4 Performance tracking

Having established the baseline GHG emissions, future GHG emission modeling for Bayer's customers will be compared to the baseline. The next steps are to estimate the specific GHG emissions based on the data to be collected by Kynetec biennially (every two years) up to 2030. BCS plans to continuously purchase

data from Kynetec to determine our progress for the achievement of our set target (see Figure 9 below for more details).

Figure 9 - Baseline establishment and performance tracking



4.2 Discussion

In this section, we describe the key findings in our assessment. The number of surveyed customers ranged from 32 (Australia cotton market) to 611 (Brazil corn market), with most markets consisting of 120-300 surveyed customers (Argentina corn, USA cotton, USA spring wheat, Canada spring wheat, Spain corn, Argentina soy, Italy corn, US winter wheat, Australia winter wheat, Mexico corn, Canada spring rape, India rice, and France winter wheat) or 500-611 surveyed customers (Brazil soybean and corn, and USA soybean and corn). The Baseline specific GHG emission varied across the 18 CCCs. Corn (0.07-0.18 kg CO₂e per kg crop in Argentina, Brazil, Italy, Spain and US) had smallest specific emissions compared to other crops, expect Mexico corn (0.42 kg CO₂e per kg crop). While cotton had higher specific emissions (0.56 kg CO₂e per kg crop in Australia and USA). Rice in India had the largest specific emission (0.97 kg CO₂e per kg crop).

The emission sources contributing to the GHG emissions varied greatly. Fertilizer decomposition was a major contributor to GHG emissions across the key markets, except for India rice. In the India rice market, methane emissions from the paddy play a major role in GHG emissions. Switching from conventional to conservation land management practices can be a GHG sink. The variance of total GHG emissions in the baseline across the 18 markets was dependent on the market share assessed, specific GHG emissions, and the production amount. The weighted specific GHG emission across 18 key markets for the baseline is **0.443 kg CO₂e per kg crop**.

Following the definition of a baseline upon which our improvement and progress will be tracked, future GHG emission modeling for Bayer's customers will be compared to the baseline specific GHG emission. We will estimate the specific GHG emissions for the 18 CCCs based on the data to be collected in the next every two or three years by Kynetec up to 2030. We will assess our 30% reduction commitment against these data estimated for baseline. Data sources in the future can be derived from internal data through Market Development trials, Carbon Business programs, or external data from similar surveys conducted in baseline.

4.3 Uncertainty analysis discussion in extant literature

In the assessment of GHG emissions, uncertainty evolves from three sources: Uncertainties on activity data (inventory), uncertainty resulting from year-to-year variability (i.e., changes in climate and management practice), and uncertainty resulting emission factors (i.e., characterization; Gibbons et al (2006).

- Uncertainty arising from inventory data can be controlled by avoiding under-representation. At the farm scale, only a little uncertainty relates to the inventory data, as data are provided directly by farmers. At landscape or regional scale, data are often based on statistical averages or expert knowledge, thus, the degree of uncertainties are typically higher compared to farm scale (Colomb, et al., 2012). Therefore, BCS has decided to partner with Kynetec to collect primary data based on interviews with farmers to ensure high accuracy of all reported activities especially those with strong influence on results, such as amount of N fertilizers reported.
- Uncertainty resulting from year-to-year variation can be reduced by using average climatic data and management practices on a several years period. For example, the same quantity of Nitrogen will result in different nitrification-denitrification rates due to variation in climatic condition (Colomb, et al., 2012). Since BCS just started the ongoing data collection in 2021 for baseline calculations, multi-year data are not yet available, but planned to be included in future to avoid such uncertainty.
- Uncertainty resulting from emission factors are associated with the chosen GHG emission calculators. Specifically, for the CFT, Clavreul et al (2017) found that the influence of model uncertainties on the GHG results are low.

4.4 Sensitivity analysis discussion in extant literature

Since data collection is still partially ongoing (expected to be complete in Spring 2022), a sensitivity analysis of the specific BCS GHG results does not exist yet. Therefore, for now, this report only provides sensitivity analysis insights on the CFT GHG calculation based on existing literature.

In a CFT case study example on the carbon footprint of open-field tomato production from 198 farms, Clavreul et al. (2017) found that several factors contribute to the variability in the carbon footprint results from CFT GHG calculation. Using a one-factor-at-a-time technique and Monte Carlo simulations, they conducted a sensitivity analysis to understand the impact of the different input parameters (farmer's inputs and model parameters) on the CFT GHG emissions results.

The results showed that the variability of total GHG emissions per ton of tomato produced was highly sensitive to variations in the production yield. Clavreul et al (2017) stated that a 70% reduction in yield resulted in a threefold increase in the GHG emission per ton of tomato. Furthermore, GHG emissions results were discovered to be sensitive towards variability in farm practices (underlined in Figure 10 below); in particular, to the ones related to fertilizer and diesel uses (e.g., for irrigation pumping).




Figure 10 - Factors that contribute to the variability in the carbon footprint results from CFT GHG calculation (Figure from Clavreul et al. (2017))

The Figure 10 shows the total GHG emissions obtained (tomato production case) with error bars portraying the minimal and maximal GHG emissions obtained when testing minimal and maximal values for each parameter one at a time. Underlined are farmer related input data. The others are model parameters.

In a very recent study, Lam et al (2021) used the CFT to evaluate possible sources of variability in GHG footprint (in terms of kg CO₂-eq/kg crop produced) of 26 crops using data from 4565 farms in 36 countries from 2013 through 2016. Across all crops and countries, they found that fertilizer use was the most important source of GHG emissions. Furthermore, they found negative relationships between GHG footprints and yields for the vast majority of the crops, suggesting that an increase in yield e.g., by growing more productive crop varieties) typically results in lower GHG footprints. According to the researchers, the reduction of GHG footprints with yield reflects that yield increase measures do not typically lead to a proportional increase in emissions. The researchers state that increases in yield are typically obtained through an increased farming efficiency which in turn does not increase GHG emission. An example is by synchronizing fertilizer application with crop nutrient requirements or by adopting more efficient crop varieties.

However, Lam et al (2021) also found several non-linear negative relationships between GHG footprints and yields for certain crops in their dataset, suggesting that optimum yield values may exist in terms of GHG footprints. For example, the GHG footprints of parsley and strawberry decreased with increasing yield, up to a certain yield value and then increased again. Therefore, several GHG improvement levers (along with yield increase) should be implemented in an orchestrated and coordinated way (Lam, et al., 2021).

- For example, with precision farming that seeks to optimize amounts, types, methods and timing of fertilizer application, yields can be increased while limiting or reducing GHG emissions from the production and application of synthetic nitrogen fertilizers
- Other opportunities to reduce GHG emissions without reducing yields are efficiency improvements of electricity and fossil fuel (e.g., by replacing inefficient machinery or substituting fossil energy)
- GHG emissions caused by electricity use for irrigation can be reduced by optimizing the efficiency of the irrigation technologies and strategies or transitioning to alternative electricity sources such as solar power.

5 Main limitations of the assessment

Relating to the limitation of the emission calculation using the CFT, the tool only considers seed emissions from potatoes and not for other crops. These could lead to an underestimation of emissions. However these emissions are reported to be quite low, compared to the other sources of emissions. The CFT plans to include this emission category in future.

In relation to Land use change (LUC), BCS acknowledges that LUC is one of the biggest contributors of GHG emissions in the global food systems. However, LUC emissions are not covered in this report due to the lack of reliable data and estimation difficulties. Therefore, BCS only included emissions which can be reliably measured in the scope of its GHG commitment. Regarding the exclusion of the production of crop protection products and fertilizers, and transportation, this is considered out of scope because the assessment focusses on emissions resulting from operations on the field.

6 Further developments of this report

In addition, the following sections will be further developed for the purpose of the further review cycles:

- Sensitivity analyses
- Uncertainty analysis
- Recommendations for further developments
- Review comments and practitioner responses
- Review report.

7. References

- Affairs, U. N. (2021, November). Sustainable Development Goals. Retrieved from The 17 Goals: <https://sdgs.un.org/goals>
- Akiyama, Hiroko, Xiaoyuan, Y., Yagi, & Kazuyuki. (2010). Evaluation of effectiveness of enhanced efficiency fertilizers as mitigation options for N₂O and NO emissions from agricultural soils: meta analysis. *Global change biology*.
- Audsley, E. (1997). Harmonisation of environmental life cycle assessment for agriculture. *The International Journal of Life Cycle Assessment*, 139.
- Bouwman, A. F., Boumans, L. J., & Batjes, N. H. (2002). Modelling global annual N₂O and NO emissions from fertilized fields. *Global Biogeochemical Cycles*, 16(4), 1080.
- Clavreul, J., Butnar, I., Rubio, V., & King, H. (2017). Intra- and inter-year variability of agricultural carbon footprints – A case study on field-grown tomatoes,. *Journal of Cleaner Production*, 158, 156-164. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0959652617309216>
- Colomb, V., Bernoux, M., Bockel, L., Chotte, J.-L., Martin, S., Martin-Phipps, C., . . . Touchemoulin, O. (2012). Review of GHG calculators in agriculture and forestry sectors - A guideline for appropriate choice and use of landscape based tools. Retrieved from https://www.fao.org/fileadmin/templates/ex_act/pdf/ADEME/Review_existingGHGtool_VF_UK4.pdf
- Cool Farm Alliance. (2020, November 13). Which IPCC Tiers does the Cool Farm Tool incorporate? Do you consider Global Warming Potential from the IPCC Assessment Report 4 or 5? Retrieved from News&Resources: <https://coolfarmtool.org/faqs/does-the-cool-farm-tool-use-global-warming-potentials-from-the-ipcc-assessment-report-4-or-ipcc-assessment-report-5/>

- Cool Farm Alliance. (2021, November). Frequently Asked Questions (FAQ). Retrieved from <https://coolfarmtool.org/coolfarmtool/frequently-asked-questions/>
- Cool Farm Alliance. (2021, November). Greenhouse gases. Retrieved from <https://coolfarmtool.org/coolfarmtool/greenhouse-gases/>
- Cool Farm Alliance. (2021, November). What reference data does the CFT use for grid electricity emission factors? Retrieved from News&Resources: <https://coolfarmtool.org/faqs/what-reference-data-does-the-cft-use-for-grid-electricity-emission-factors/>
- Cool Farm Alliance. (2022). Cool Farm Tool: Updates to the 2019 IPCC Guidelines for Greenhouse Gas Inventories.
- EC, E. C., JRC, J., & IES, I. (2010). International Reference Life Cycle Data System (ILCD) Handbook - General Guide for Life Cycle Assessment - Detailed guidance. Luxembourg: Publications Office of the European Union.
- FAO. (2010). Global Ecological Zones for FAO Forest Reporting: 2010 Update. Forest Resources Assessment Working Paper 179.
- FAO. (2021). The share of food systems in total greenhouse gas emissions. Global, regional and country trends, 1990–2019. FAOSTAT Analytical Brief Series No. 31. Rome.
- Fatrelli, P. (2017). Heat energy saving of grain dryers.
- GHGprotocol. (2003). Emission factors from cross sector tools. URL - <http://www.ghgprotocol.org/calculation-tools/all-tools>.
- Gibbons, J. M., Ramsden, S. J., & Blake, A. (2006). Modelling uncertainty in greenhouse gas emissions from UK agriculture at the farm level. *Agriculture, Ecosystems and Environment*, 112(4), 347-355. doi:<https://doi.org/10.1016/j.agee.2005.08.029>
- Hillier, J., Walter, C., Malin, D., Garcia-Suarez, T., Mila-i-Canals, L., & Smith, P. (2011). A farm-focused calculator for emissions from crop and livestock production. *Environmental Modelling & Software*, 26(9), 1070-1078. Retrieved from <https://www.sciencedirect.com/science/article/pii/S1364815211000892>
- IFA and FAO. (2001). Global estimates of gaseous emissions of NH₃, NO and N₂O from agricultural land. Food and Agriculture Organization of the United Nations (FAO).
- International Organization for Standardization (ISO). (2006). ISO 14040. Environmental management - Life cycle assessment - Principles and framework, 23. Geneva.
- IPCC. (2019). Climate Change and Land - An IPCC Special Report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems. Retrieved from <https://www.ipcc.ch/site/assets/uploads/sites/4/2021/07/210714-IPCCJ7230-SRCCL-Complete-BOOK-HRES.pdf>
- Khaledian, M. M. (2014). Diesel oil consumption, work duration, and crop production of corn and durum wheat under conventional and no-tillage in southeastern France. *Archives of Agronomy and Soil Science*.
- Kynetec. (2021, 11 19). FarmTrak. Retrieved from kynetec: <https://www.kynetec.com/farmtrak-agriculture-tracking-studies>
- kynetec. (2021, November). Tracking Studies. Retrieved from <https://www.kynetec.com/farmtrak-agriculture-tracking-studies>
- Lam, W. Y., Sim, S., Kulak, M., van Zelm, R., Schipper, A. M., & Huijbregts, M. A. (2021). Drivers of variability in greenhouse gas footprints of crop production. *Journal of Cleaner Production*, 315. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0959652621023398>
- Ogle S.M., B. F. (2005). . Agricultural management impacts on soil organic carbon storage under moist and dry climatic conditions of temperate and tropical regions. *Biogeochemistry*, 1, pp. 87 - 121.
- Sadaka, S. (2014). On-Farm drying and storage of Soybeans. Soybean production handbook. University of Arkansas System, Division of Agriculture.
- Science Based Targets. (2021). Science Based Targets - How it works. Retrieved from <https://sciencebasedtargets.org/how-it-works>

- Science-Based Targets Initiative. (2021, November). Retrieved from How-it-works:
<https://sciencebasedtargets.org/how-it-works>
- Smith P., P. D. (1997). Potential for carbon sequestration in european soils: Preliminary estimates for five scenarios using results from long-term experiments. *Global Change Biology*, 1, pp. 67 - 79.
- Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Ingo, F., M., B. E., . . . R., C. S. (2015). Planetary boundaries: Guiding human development on a changing planet. American Association for the Advancement of Science.
- UN, United Nations. (2021, 11 19). Sustainable Development Goals, SDGs. Retrieved from The 17 Goals:
<https://sdgs.un.org/goals>
- WRI, World Resources Institute. (2004). The Greenhouse Gas Protocol. Retrieved from GHG Protocol, p37: <https://ghgprotocol.org/sites/default/files/standards/ghg-protocol-revised.pdf>
- Xiaoyuan Yan, K. Y. (2005). Statistical analysis of the major variables controlling methane emission from rice fields. *Global Change Biology*, 11(7), pp. 1131-1141. doi: doi: 10.1111/j.1365-2486.2005.00976.x.

8. Appendix

Table 19 - Summary of specific GHG emission (kg CO₂e per kg crop) from surveyed BCS customer farms by gate-to-gate emission source in the baseline year for the 18 CCC's.

CCC	CFT modeled GHG Emission (kg CO ₂ e)	Total Crop Weight of Surveyed Customer (tonnes)	Specific GHG Emission (kg CO ₂ e per kg crop)	Distribution of Specific GHG Emission per Emission sources (kg CO ₂ e per kg crop)					
				Fertilizer/ Soil decomposition	Irrigation	Machinery	Management changes	Crop Residue	Paddy methane
Argentina-corn	148839197	1109361.9	0.1342	0.0956	0	0.0191	-0.0123	0.0318	-
Australia-cotton	5236418.46	9408.22	0.5566	0.382	0.0441	0.0528	0.0134	0.0642	-
Brazil-corn	135548634.1	774304.77	0.1751	0.178	0.0002	0.0149	-0.0458	0.0278	-
Brazil-soybean	150393044.6	584936.18	0.2571	0.2606	0.0011	0.0328	-0.0826	0.0453	-
India-rice, paddy	3526317.05	3634.71	0.9702	0.2249	0.0435	0.0352	0.0008	0.071	0.5948
Italy-corn	4612713.74	64558.41	0.0715	0.0397	0.0075	0.0068	-0.0059	0.0233	-
Spain-corn	10641954.47	95512.85	0.1114	0.0635	0.0196	0.0091	-0.0053	0.0246	-
USA-corn	272990301.3	2909794.63	0.0938	0.055	0.0053	0.0116	-0.0058	0.0278	-
USA-cotton	49354335.23	88910.75	0.5551	0.3228	0.0673	0.0954	-0.0158	0.0854	-
USA-soybean	94663958.61	683436.8	0.1385	0.0755	0.0107	0.0349	-0.0318	0.0493	-
Argentina-soybean	130193457.4	344495.7	0.3779	0.293	0	0.039	-0.0071	0.0531	-
Australia-wheat	4775618.29	7877.66	0.6062	0.5335	0	0.0497	-0.0255	0.0485	-
Canada-rapeseed	33124891.18	75533.29	0.4385	0.3281	0	0.0637	-0.0099	0.0567	-
Canada-wheat	19814089.57	61022.7	0.3247	0.23	0.0017	0.0488	-0.0026	0.0468	-
France-wheat	6221656.53	32100.5	0.1938	0.1385	0.0008	0.0234	0.0013	0.0298	-
Mexico-corn	26203826.23	62709.07	0.4179	0.3671	0.0131	0.0171	-0.0013	0.0219	-
USA-wheat, Spring	13184910.7	45565.18	0.2894	0.1926	0.0226	0.0492	-0.018	0.043	-
USA-wheat, Winter	13481567.08	54669.41	0.2466	0.1687	0.0155	0.0348	-0.0173	0.0449	-