



Document Title

**Summary of the residues in or on treated products, food and feed for
fosetyl-aluminium + fluopicolide WG 71.11 (666.7 + 44.4 g/kg)**

Data Requirements

EU Regulation 1107/2009 & EU Regulation 284/2013

Document MCP

Section 8: Residues in or on treated products, food and feed

According to the Guidance Document SANCO/10181/2013 for
preparing dossiers for the approval of a chemical active substance

Date

2015-09-01

Author(s)

Bayer CropScience



M-532104-01-2

*This document is copyright protected.
Any distribution, reproduction or publication requires
the consent of Bayer AG (or its respective affiliate).
Any use of the document or its content for regulatory or
any other commercial purpose is prohibited and constitutes
a violation of the underlying license agreement.*



OWNERSHIP STATEMENT

This document, the data contained in it and copyright therein are owned by Bayer CropScience. No part of the document or any information contained therein may be disclosed to any third party without the prior written authorisation of Bayer CropScience.

The summaries and evaluations contained in this document are based on unpublished proprietary data submitted for the purpose of the assessment undertaken by the regulatory authority. Other registration authorities should not grant, amend, or renew a registration on the basis of the summaries and evaluation of unpublished proprietary data contained in this document unless they have received the data on which the summaries and evaluation are based, either:

- from Bayer CropScience; or
- from other applicants once the period of data protection has expired.

This document is copyright protected. Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliates). Any use of the document or its content for any other commercial purpose is prohibited and constitutes a violation of the underlying license agreement.



Document MCP – Section 8: Residues in or on treated products, food and feed
Fosetyl-aluminium + fluopicolide WG 71.11

Version history

Date	Data points containing amendments or additions ¹ and brief description	Document identifier and version number

¹ It is suggested that applicants adopt a similar approach to showing revisions and version history as outlined in SANCO/10180/2013 Chapter 4 How to revise an Assessment Report

This document is copyright protected (equivalent to a trademark) and its distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliate) or any other commercial purpose is prohibited for regulatory or a violation of the underlying license agreement.



Table of Contents

CP 8	RESIDUES IN OR ON TREATED PRODUCTS, FOOD OR FEED	Page 5
------	--	-----------

*This document is copyright protected.
 Any distribution, reproduction or publication requires
 the consent of Bayer AG (or its respective affiliate).
 Any use of the document or its content for regulatory or
 any other commercial purpose is prohibited and constitutes
 a violation of the underlying license agreement.*



Document MCP – Section 8: Residues in or on treated products, food and feed
Fosetyl-aluminium + fluopicolide WG 71.11

CP 8 RESIDUES IN OR ON TREATED PRODUCTS, FOOD OR FEED

Fosetyl was included in Annex I to Directive 91/414/EEC in 2006 (Directive 2006/64/CE of 18 July 2006, Entry into Force on 1 May 2007). Fosetyl is the ISO common name for ethyl hydrogen phosphonate (IUPAC) but the aluminium salt fosetyl-aluminium (fosetyl-Al), a variant of fosetyl, is used in the formulated product.

The representative formulation fosetyl-Al + fluopicolide WG 71.11 is a water-dispersible granules (WG) formulation containing 666.7 g/kg of fosetyl-Al and 44.4 g/kg of fluopicolide. The use of fosetyl-Al + fluopicolide WG 71.11 on grape against *Plasmopara viticola* has been selected as representative use for the approval renewal of fosetyl. The critical Good Agricultural Practice (cGAP) supported at the European level in the approval renewal process of fosetyl consists of 2 foliar spray applications between BBCH 15 and 81 at 2.0 kg/ha of fosetyl-Al with an interval between applications of 10 to 14 days and a PHI of 21 days. All relevant metabolism and residue data in support of this use are summarized in Document MCA, Section 6.

This document is copyright protected and requires the consent of Bayer AG (or its respective affiliates) for any distribution, reproduction or publication. Any use of the document or its content for regulatory or commercial purposes is prohibited and constitutes a violation of the underlying license agreement.