





## Appendix III


Admire PRO label highlighting  
specific precautions to protect bees





|  |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
|--|--|---------------------|--|-------------------------------|--|-------------------|---|-------------------|---|---|--|--|--|
| 3.875"   |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| 5.5"   |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| <p style="text-align: center;"><b>FIRST AID</b></p> <table border="1"> <tr> <td><b>IF SWALLOWED</b></td><td> <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> </td></tr> <tr> <td><b>IF ON SKIN OR CLOTHING</b></td><td> <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul> </td></tr> <tr> <td><b>IF INHALED</b></td><td> <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul> </td></tr> <tr> <td><b>IF IN EYES</b></td><td> <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul> </td></tr> <tr> <td colspan="2"> <p style="text-align: center;"><b>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</b></p> </td></tr> <tr> <td colspan="2"> <p><b>Note To Physician:</b> No specific antidote is available. Treat the patient symptomatically.</p> </td></tr> </table> |  | <b>IF SWALLOWED</b> | <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> | <b>IF ON SKIN OR CLOTHING</b> | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul> | <b>IF INHALED</b> | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul> | <b>IF IN EYES</b> | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul> | <p style="text-align: center;"><b>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</b></p> |  | <p><b>Note To Physician:</b> No specific antidote is available. Treat the patient symptomatically.</p> |  |
| <b>IF SWALLOWED</b>  | <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| <b>IF ON SKIN OR CLOTHING</b>  | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>   |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| <b>IF INHALED</b>  | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| <b>IF IN EYES</b>  | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| <p style="text-align: center;"><b>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</b></p>  |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| <p><b>Note To Physician:</b> No specific antidote is available. Treat the patient symptomatically.</p>   |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| 5.0"   |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| <p style="text-align: center;"><b>PRECAUTIONARY STATEMENTS</b></p> <p><b>HAZARDS TO HUMANS AND DOMESTIC ANIMALS</b></p> <p><b>CAUTION</b></p> <p>Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.</p> <p><b>Applicators and Other Handlers Must Wear:</b></p> <ul style="list-style-type: none"> <li>• Long-sleeved shirt and long pants</li> <li>• Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.</li> <li>• Shoes plus socks</li> </ul> <p>Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.</p> <p><b>ENGINEERING CONTROLS STATEMENTS</b></p> <ul style="list-style-type: none"> <li>• When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.</li> </ul>  |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| 1  |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |
| 4.375"   |  |                     |  |                               |  |                   |   |                   |   |   |  |  |  |











#### Application Directions (Soil)

Direct applications of ADMIRE PRO SYSTEMIC PROTECTANT into the seed or root-zone of crop. Failure to place ADMIRE PRO SYSTEMIC PROTECTANT into root-zone may result in loss of control or delay in onset of activity. Apply ADMIRE PRO SYSTEMIC PROTECTANT by ground application or chemigation application. For seedling flats or trays, only apply with broadcast, foliar applications or where product is intended to be washed from foliage to soil prior to drying on foliage.

Optimum activity of ADMIRE PRO SYSTEMIC PROTECTANT results from applications to the root-zone of plants to be protected. The earlier ADMIRE PRO SYSTEMIC PROTECTANT is available to a developing plant, the earlier the protection begins. ADMIRE PRO SYSTEMIC PROTECTANT is continuously taken into the roots over a long period of time and the systemic nature of ADMIRE PRO SYSTEMIC PROTECTANT allows movement from roots through the xylem tissue to all vegetative parts of the plant. This results in residual activity of ADMIRE PRO

(continued)




← 3.875 →

**5.5"**

**5.0"**

7

← 4.375" →



3.875"

**5.5"**

A vertical dimension line with arrows at both ends, indicating a height of 5.0 inches between two horizontal lines.

**5.5"**

**5.0"**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (**RPZ**) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the **RPZ**, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check

9

← 4.375" →

3.875"

valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

#### ROTATIONAL CROPS\*

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.

#### IMMEDIATE PLANT-BACK:

All crops on this label plus the following crops not on this label: barley, canola, corn (field, pop & sweet), rapeseed, sorghum, and wheat.

#### 30-DAY PLANT-BACK:

Cereals (including buckwheat, millet, oats, rice, rye, and triticale), safflower

#### 12-MONTH PLANT-BACK:

All Other Crops

\*Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

#### FIELD CROPS

#### Application Rates – ADMIRE PRO Systemic Protectant

#### COTTON – SOIL

| Pests Controlled |            | Rate<br>fluid ounces/Acre               |
|------------------|------------|---|
| Cotton aphid     | Thrips     | 7.4 – 9.2<br>(Depending on row-spacing) |
| Plant bugs       | Whiteflies |   |

#### Cotton – Soil Applications

Apply specified dosage in one of the following methods:

1. In-furrow spray during planting directed on or below seed;
2. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting;
3. Chemigation into root-zone through low-pressure drip or trickle irrigation.

(continued)

4.375"

| 3.875"  |   |                  |                           |   |           |                         |  |  |
|---|---|------------------|---------------------------|---|-----------|-------------------------|--|--|
| 5.5"  | <p><b>COTTON – SOIL (continued)</b></p> <p><b>Cotton – Soil Application Restrictions</b><br/> Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>9.2 fluid ounces/Acre</b> (0.33 lb AI/Acre)<br/> Regardless of formulation or method of application, apply no more than 0.5 lb active ingredient per acre per year, including seed treatment, soil and foliar uses. Do not graze treated fields after any application of ADMIRE PRO SYSTEMIC PROTECTANT. Please see Resistance Management section of this label.</p>  |                  |                           |   |           |                         |  |  |
|   | <p><b>COTTON – FOLIAR</b></p> <table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Cotton aphid<br/>Cotton fleahopper<br/>Bandedwinged whitefly<br/>Plant bugs (excludes <i>Lygus hesperus</i>)<br/>Green stink bug<br/>Southern green stink bug<br/>Bollworm/Budworm (ovicidal effect)</td><td>0.9 – 1.7</td></tr> <tr> <td colspan="2"><b>Pests Suppressed</b></td></tr> <tr> <td>Lygus bug (<i>Lygus hesperus</i>)<br/>Whiteflies (other than Bandedwinged whitefly)</td><td>1.3 – 1.7</td></tr> </table> <p><b>Cotton – Foliar Applications</b><br/> Apply ADMIRE PRO SYSTEMIC PROTECTANT through properly calibrated ground or aerial application equipment.</p> <p><b>Cotton – Foliar Application Restrictions</b><br/> Pre-Harvest Interval (PHI): <b>14 days</b><br/> Minimum interval between foliar applications: <b>7 days</b><br/> Maximum foliar applied ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>8.7 fluid ounces/Acre</b> (0.31 lb AI/Acre)<br/> Regardless of formulation or method of application, apply no more than 0.5 lb active ingredient per acre per year, including seed treatment, soil and foliar uses. Do not graze treated fields after any application of ADMIRE PRO SYSTEMIC PROTECTANT. Please see Resistance Management section of this label.</p> | Pests Controlled | Rate<br>fluid ounces/Acre | Cotton aphid<br>Cotton fleahopper<br>Bandedwinged whitefly<br>Plant bugs (excludes <i>Lygus hesperus</i> )<br>Green stink bug<br>Southern green stink bug<br>Bollworm/Budworm (ovicidal effect) | 0.9 – 1.7 | <b>Pests Suppressed</b> |  | Lygus bug ( <i>Lygus hesperus</i> )<br>Whiteflies (other than Bandedwinged whitefly) |
| Pests Controlled  | Rate<br>fluid ounces/Acre   |                  |                           |   |           |                         |  |  |
| Cotton aphid<br>Cotton fleahopper<br>Bandedwinged whitefly<br>Plant bugs (excludes <i>Lygus hesperus</i> )<br>Green stink bug<br>Southern green stink bug<br>Bollworm/Budworm (ovicidal effect) | 0.9 – 1.7   |                  |                           |   |           |                         |  |  |
| <b>Pests Suppressed</b>   |   |                  |                           |   |           |                         |  |  |
| Lygus bug ( <i>Lygus hesperus</i> )<br>Whiteflies (other than Bandedwinged whitefly)  | 1.3 – 1.7   |                  |                           |   |           |                         |  |  |
| 4.375"  |   |                  |                           |   |           |                         |  |  |



| 3.875"   |  |                  |                           |   |           |                                    |  |   |
|--|--|------------------|---------------------------|---|-----------|------------------------------------|--|---|
| 5.5"   | <b>PEANUT – FOLIAR <sup>1/</sup> (continued)</b><br><b>Peanut – Foliar Application Restrictions</b><br>Pre-Harvest Interval (PHI): <b>14 days</b><br>Maximum foliar applied ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>3.7 fluid ounces/Acre</b><br>(0.13 lb AI/Acre)<br>Minimum interval between foliar applications: <b>5 days</b><br><sup>1/</sup> Use not permitted in California unless otherwise directed by state specific 24(c) labeling.   |                  |                           |   |           |                                    |  |   |
|  | <b>POTATO – SOIL</b><br><table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Aphids<br/>Colorado potato beetle<br/>Flea beetles<br/>Leafhoppers<br/>Potato psyllid</td><td>5.7 – 8.7</td></tr> <tr> <td colspan="2"><b>Pests / Diseases Suppressed</b></td></tr> <tr> <td>Symptoms of:<br/>Potato leaf roll virus (PLRV)<br/>Potato yellows<br/>Net necrosis<br/>Wireworms (with in-furrow spray at-planting)</td><td>5.7 – 8.7</td></tr> </table> <b>Potato – Soil Applications</b><br>Apply specified dosage in one of the following methods:<br>1. In-furrow spray during planting directed on seed pieces or seed potatoes;<br>2. Subsurface side-dress on both sides of the row covered with 3 or more inches of soil;<br>3. Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil;<br>4. Narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting. For effective pest control or suppression, ADMIRE PRO SYSTEMIC PROTECTANT applications must be placed below soil-surface and in contact with seed piece or within root-zone. For potatoes grown on highly permeable soils with shallow water table, at-plant applications of ADMIRE PRO SYSTEMIC PROTECTANT may be made in a 2 to 4 inch band (width of planter shoe opening) and completely covered. | Pests Controlled | Rate<br>fluid ounces/Acre | Aphids<br>Colorado potato beetle<br>Flea beetles<br>Leafhoppers<br>Potato psyllid | 5.7 – 8.7 | <b>Pests / Diseases Suppressed</b> |  | Symptoms of:<br>Potato leaf roll virus (PLRV)<br>Potato yellows<br>Net necrosis<br>Wireworms (with in-furrow spray at-planting) |
| Pests Controlled   | Rate<br>fluid ounces/Acre  |                  |                           |   |           |                                    |  |   |
| Aphids<br>Colorado potato beetle<br>Flea beetles<br>Leafhoppers<br>Potato psyllid  | 5.7 – 8.7  |                  |                           |   |           |                                    |  |   |
| <b>Pests / Diseases Suppressed</b>   |  |                  |                           |   |           |                                    |  |   |
| Symptoms of:<br>Potato leaf roll virus (PLRV)<br>Potato yellows<br>Net necrosis<br>Wireworms (with in-furrow spray at-planting)                          | 5.7 – 8.7  |                  |                           |   |           |                                    |  |   |
| <b>Potato – Soil Application Restrictions</b><br>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>8.7 fluid ounces/Acre</b> (0.31 lb AI/Acre) |  |                  |                           |   |           |                                    |  |   |
| 13   |  |                  |                           |   |           |                                    |  |   |
| 4.375"   |  |                  |                           |   |           |                                    |  |   |
| 5.0"   |  |                  |                           |   |           |                                    |  |   |

3.875"

5"

| <b>Pests Controlled</b>   | <b>Rate<br/>fluid ounces/100 lbs seed</b> | <b>Rate<br/>fluid ounces/Acre*</b> |
|---|---|------------------------------------|
| Aphids<br>Colorado potato beetle<br>Flea beetles<br>Leafhoppers<br>Potato psyllid<br>Wireworms<br>(seed-piece protection)   | 0.17 – 0.35                               | 3.5 – 7.0                          |
| <b>Diseases Suppressed</b>  |   |                                    |
| Symptoms of:<br>Potato leaf roll virus (PLRV)<br>Potato yellows<br>Net necrosis   | 0.35                                      | 7.0                                |
| <b>Potato – Seed-piece Application</b>  |   |                                    |
| Apply specified dosage as a diluted spray onto seed-pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part ADMIRE PRO SYSTEMIC PROTECTANT. Agitate or stir spray solution as needed. Fungicidal or inert absorbent dusts may be applied after ADMIRE PRO SYSTEMIC PROTECTANT application. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed-pieces as soon as possible after treating avoiding prolonged exposure of ADMIRE PRO SYSTEMIC PROTECTANT treated seed-pieces to sunlight and in accordance with the recommendation of your local Extension specialist. |   |                                    |
| <b>Potato – Seed-piece Application Restrictions</b>   |   |                                    |
| Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>8.7 fluid ounces/Acre</b> (0.31 lb AI/Acre)   |   |                                    |
| Do not use treated seed-pieces for food, feed, or fodder.   |   |                                    |
| * BASED ON A SEEDING RATE OF 2000 LBS/ACRE.   |   |                                    |

$O''$


| <b>Pests Controlled</b>  | <b>Rate<br/>fluid ounces/Acre</b> |
|--|-----------------------------------|
| Aphids                      Leafhoppers<br>Colorado potato beetle    Potato psyllid<br>Flea beetles  | 1.3                               |
| <b>Potato – Foliar Applications</b><br>Apply ADMIRE PRO SYSTEMIC PROTECTANT through properly calibrated ground and aerial application equipment. |                                   |

14

4.375"







| 3.875"   |                           |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
|--|---------------------------|------------------|---------------------------|---|------------|---------------------|--|--|------------|---|--|---|--|--|--|
| <p><b>VEGETABLE and SMALL FRUIT CROPS</b><br/> <b>Application Rates – ADMIRE PRO Systemic Protectant</b></p> <p><b>CUCURBIT VEGETABLES – SOIL <sup>1/</sup></b><br/> <b>Crops of Crop Group 9 including:</b> Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of <i>Cucumis melo</i> including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of <i>Citrullus lanatus</i>)</p> <table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Aphids<br/>Cucumber beetles<br/>Leafhoppers<br/>Thrips (foliage feeding thrips only)<br/>Whiteflies</td><td>7.0 – 10.5</td></tr> <tr> <th colspan="2">Diseases Suppressed</th></tr> <tr> <td>Bacterial wilt (as vectored by various cucumber beetles)<br/>Leaf silvering resulting from whitefly feeding</td><td>7.0 – 10.5</td></tr> <tr> <td colspan="2"> <b>Cucurbit Vegetables – Soil Applications</b><br/>           Apply specified dosage in one of the following methods:<br/>           1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;<br/>           2. In-furrow spray directed on or below seed;<br/>           3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application;<br/>           4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;<br/>           5. Post-seeding drench, transplant-water drench, or hill drench;<br/>           6. Subsurface side-dress on both sides of each row. ADMIRE PRO SYSTEMIC PROTECTANT must be incorporated into root-zone.         </td></tr> <tr> <td colspan="2"> <b>Cucurbit Vegetables – Soil Application Restrictions</b><br/>           Pre-Harvest Interval (PHI): <b>21 days</b><br/>           Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season: <b>10.5 fluid ounces/Acre</b> (0.38 lb AI/Acre)         </td></tr> <tr> <td colspan="2"><sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.</td></tr> </table> |                           | Pests Controlled | Rate<br>fluid ounces/Acre | Aphids<br>Cucumber beetles<br>Leafhoppers<br>Thrips (foliage feeding thrips only)<br>Whiteflies | 7.0 – 10.5 | Diseases Suppressed |  | Bacterial wilt (as vectored by various cucumber beetles)<br>Leaf silvering resulting from whitefly feeding | 7.0 – 10.5 | <b>Cucurbit Vegetables – Soil Applications</b><br>Apply specified dosage in one of the following methods:<br>1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;<br>2. In-furrow spray directed on or below seed;<br>3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application;<br>4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;<br>5. Post-seeding drench, transplant-water drench, or hill drench;<br>6. Subsurface side-dress on both sides of each row. ADMIRE PRO SYSTEMIC PROTECTANT must be incorporated into root-zone. |  | <b>Cucurbit Vegetables – Soil Application Restrictions</b><br>Pre-Harvest Interval (PHI): <b>21 days</b><br>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season: <b>10.5 fluid ounces/Acre</b> (0.38 lb AI/Acre) |  | <sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling. |  |
| Pests Controlled   | Rate<br>fluid ounces/Acre |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| Aphids<br>Cucumber beetles<br>Leafhoppers<br>Thrips (foliage feeding thrips only)<br>Whiteflies  | 7.0 – 10.5                |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| Diseases Suppressed  |                           |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| Bacterial wilt (as vectored by various cucumber beetles)<br>Leaf silvering resulting from whitefly feeding   | 7.0 – 10.5                |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| <b>Cucurbit Vegetables – Soil Applications</b><br>Apply specified dosage in one of the following methods:<br>1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;<br>2. In-furrow spray directed on or below seed;<br>3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application;<br>4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;<br>5. Post-seeding drench, transplant-water drench, or hill drench;<br>6. Subsurface side-dress on both sides of each row. ADMIRE PRO SYSTEMIC PROTECTANT must be incorporated into root-zone.  |                           |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| <b>Cucurbit Vegetables – Soil Application Restrictions</b><br>Pre-Harvest Interval (PHI): <b>21 days</b><br>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season: <b>10.5 fluid ounces/Acre</b> (0.38 lb AI/Acre)  |                           |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| <sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.   |                           |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| 5.5"   |                           |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| 5.0"   |                           |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |
| 4.375"   |                           |                  |                           |   |            |                     |  |  |            |   |  |   |  |  |  |



3.875"

**BULB VEGETABLE (*Allium* sp.) Group 3-07 – SOIL <sup>1/</sup>**

**Crops of Crop Group 3-07 Including:** Chive (fresh leaves), Chinese chive (fresh leaves), Daylily (bulb), Elegans hosta, Fritillaria (bulb and leaves), Garlic (common group, great-headed group, serpent group), Kurrat group, Leek group (including common, lady's and wild), Lily (bulb), Onion (bulb and green leaves including: common group, Beltsville bunching, Chinese bulb, fresh, green, macrostem, Pearl group, potato onion group, tree onion-tops, Welsh-tops), Shallot, plus cultivars, varieties, and/or hybrids of these

| Pests Controlled  | Rate<br>fluid ounces/Acre |
|---|---------------------------|
| Thrips (foliage feeding thrips only)  | 14.0                      |
| <b>Bulb Vegetable – Soil Applications</b><br>Apply specified dosage in one of the following methods:<br>1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;<br>2. In-furrow spray directed on or below seed;<br>3. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;<br>4. Post-seeding drench, transplant-water drench, or hill drench. |                           |
| <b>Bulb Vegetable – Soil Application Restrictions</b><br>Pre-Harvest Interval (PHI): <b>21 days</b><br>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season: <b>14.0 fluid ounces/Acre</b><br>(0.5 lb AI/Acre)<br>Applications made to higher organic matter soils may result in reduced or shortened activity on pest.   |                           |
| <sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.  |                           |

**GREENHOUSE VEGETABLES – SOIL <sup>1/</sup>**

(Mature Cucumber and Tomato plants only in production greenhouses)

| Pests Controlled   | Rate<br>fluid ounces/1000 plants |
|--|----------------------------------|
| Aphids Whiteflies  | 0.6                              |
| <b>Greenhouse Vegetables – Soil Applications</b><br>Apply specified dosage in a minimum of 16 gallons of water for tomatoes and 21 gallons of water for cucumbers using soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. Apply only to plants grown in field-type soils, potting media, or mixtures thereof. Do not apply to plants grown in non-soil medias such as perlite, vermiculite, rock wool or other soil-less media, or plants growing hydroponically. Do not apply to immature plants since phytotoxicity may occur.<br>Apply when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials ( <i>Orius</i> sp.) can occur when ADMIRE PRO SYSTEMIC PROTECTANT is applied.<br>Many varieties of vegetables have been tested for tolerance to ADMIRE PRO SYSTEMIC PROTECTANT and show good safety. However, certain varieties may show more sensitivity to ADMIRE PRO SYSTEMIC PROTECTANT. Therefore, treatment of a few plants is recommended before treating the whole greenhouse. |                                  |

(continued)

4.375"

| 3.875"   |   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
|--|---|------------------|---------------------------|---|-----------------|--|------------|--|-------------|--|------------|--------------|-----------------|---------------------|------------|---------------------------|-------------|-------------------------------|
| 5.5"   | <p><b>GREENHOUSE VEGETABLES – SOIL <sup>1/</sup> (continued)</b></p> <p><b>Greenhouse Vegetables – Soil Application Restrictions</b><br/> Pre-Harvest Interval (PHI): <b>0 day</b><br/> Maximum number ADMIRE PRO SYSTEMIC PROTECTANT applications per crop season: <b>1</b>.</p> <p><sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.</p>   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
|  | <p><b>FRUITING VEGETABLES – SOIL <sup>1/</sup></b><br/> <b>Crops of Crop Group 8 plus Okra including:</b> Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet) Tomato, Pepinos, Tomatillo</p> <p><b>Field Application Rates. See details below for additional planthouse rates.</b></p> <table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Aphids                      Leafhoppers</td><td>Okra and Pepper</td></tr> <tr> <td>Colorado potato beetle      Thrips (foliage feeding thrips only)</td><td>7.0 – 14.0</td></tr> <tr> <td>Flea beetles                      Whiteflies</td><td>Other Crops</td></tr> <tr> <td></td><td>7.0 – 10.5</td></tr> </table> <p><b>Diseases Suppressed</b></p> <table> <tr> <td>Symptoms of:</td><td>Okra and Pepper</td></tr> <tr> <td>Tomato mottle virus</td><td>7.0 – 14.0</td></tr> <tr> <td>Tomato spotted wilt virus</td><td>Other Crops</td></tr> <tr> <td>Tomato yellow leaf curl virus</td><td>7.0 – 10.5</td></tr> </table> <p><b>Fruiting Vegetables – Soil Applications</b><br/> Apply specified dosage in one of the following methods:<br/> 1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;<br/> 2. In-furrow spray directed on or below seed;<br/> 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application;<br/> 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;<br/> 5. Post-seeding drench, transplant-water drench, or hill drench;<br/> 6. Subsurface side-dress on both sides of each row. ADMIRE PRO SYSTEMIC PROTECTANT must be incorporated into root-zone.</p> <p><b>Fruiting Vegetable – Soil Application Restrictions</b><br/> Pre-Harvest Interval (PHI): <b>21 days</b><br/> Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed on pepper and okra crops per crop season: <b>14.0 fluid ounces/Acre</b> (0.5 lb AI/Acre)<br/> Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed on other fruiting vegetable crops per crop season: <b>10.5 fluid ounces/Acre</b> (0.38 lb AI/Acre)<br/> <sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.</p> | Pests Controlled | Rate<br>fluid ounces/Acre | Aphids                      Leafhoppers | Okra and Pepper | Colorado potato beetle      Thrips (foliage feeding thrips only) | 7.0 – 14.0 | Flea beetles                      Whiteflies | Other Crops |  | 7.0 – 10.5 | Symptoms of: | Okra and Pepper | Tomato mottle virus | 7.0 – 14.0 | Tomato spotted wilt virus | Other Crops | Tomato yellow leaf curl virus |
| Pests Controlled   | Rate<br>fluid ounces/Acre   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| Aphids                      Leafhoppers                          | Okra and Pepper   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| Colorado potato beetle      Thrips (foliage feeding thrips only) | 7.0 – 14.0  |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| Flea beetles                      Whiteflies                     | Other Crops   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
|  | 7.0 – 10.5  |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| Symptoms of:   | Okra and Pepper   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| Tomato mottle virus  | 7.0 – 14.0  |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| Tomato spotted wilt virus  | Other Crops   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| Tomato yellow leaf curl virus                                    | 7.0 – 10.5  |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| 20   |   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |
| 4.375"   |   |                  |                           |   |                 |  |            |  |             |  |            |              |                 |                     |            |                           |             |                               |









| 3.875"  |  |                  |                           |   |     |                  |                           |   |            |
|---|--|------------------|---------------------------|---|-----|------------------|---------------------------|---|------------|
| 5.5"  | <div> <div> <b>HERBS – FOLIAR</b><br/> <b>Crops of Crop Subgroup 19A including:</b> Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood. </div> <table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Aphids<br/>Flea beetles<br/>Leafhoppers<br/>Whiteflies</td><td>1.2</td></tr> </table> <div> <b>Herbs – Foliar Applications</b><br/> Apply ADMIRE PRO SYSTEMIC PROTECTANT through properly calibrated ground and aerial application equipment. </div> <div> <b>Herbs – Foliar Application Restrictions</b><br/> Pre-Harvest Interval (PHI): <b>7 days</b><br/> Minimum interval between applications: <b>5 days</b><br/> Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per season: <b>3.6 fluid ounces/Acre</b> (0.13 lb AI/Acre) </div> <div> <b>Herbs – Foliar Application Note</b><br/> Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, Bayer CropScience strongly recommends that only small areas or numbers of plants of each be treated and evaluated prior to commercial use. </div> <div> <b>BRASSICA (COLE) LEAFY VEGETABLES – SOIL <sup>19</sup></b><br/> <b>Crops of Crop Group 5 including:</b> Broccoli, Broccoli raab (<i>rapini</i>), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (<i>gai lan</i>) broccoli, Chinese (<i>bok choy</i>) cabbage, Chinese (<i>napa</i>) cabbage, Chinese mustard (<i>gai choy</i>) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens </div> <table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Aphids<br/>Leafhoppers<br/>Thrips (foliage feeding thrips only)<br/>Whiteflies</td><td>4.4 – 10.5</td></tr> </table> <div>(continued)</div> </div> | Pests Controlled | Rate<br>fluid ounces/Acre | Aphids<br>Flea beetles<br>Leafhoppers<br>Whiteflies | 1.2 | Pests Controlled | Rate<br>fluid ounces/Acre | Aphids<br>Leafhoppers<br>Thrips (foliage feeding thrips only)<br>Whiteflies | 4.4 – 10.5 |
| Pests Controlled  | Rate<br>fluid ounces/Acre  |                  |                           |   |     |                  |                           |   |            |
| Aphids<br>Flea beetles<br>Leafhoppers<br>Whiteflies                         | 1.2  |                  |                           |   |     |                  |                           |   |            |
| Pests Controlled  | Rate<br>fluid ounces/Acre  |                  |                           |   |     |                  |                           |   |            |
| Aphids<br>Leafhoppers<br>Thrips (foliage feeding thrips only)<br>Whiteflies | 4.4 – 10.5   |                  |                           |   |     |                  |                           |   |            |
| 5.0"  | 24   |                  |                           |   |     |                  |                           |   |            |
| 4.375"  |  |                  |                           |   |     |                  |                           |   |            |

| 3.875"  |  |                  |                           |   |
|---|--|------------------|---------------------------|---|
| 5.5"  | <p><b>BRASSICA (COLE) LEAFY VEGETABLES – SOIL <sup>1/</sup> (continued)</b></p> <p><b>Brassica Leafy Vegetables – Soil Applications</b><br/>           Apply specified dosage in one of the following methods:</p> <ol style="list-style-type: none"> <li>1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;</li> <li>2. In-furrow spray directed on or below seed;</li> <li>3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application;</li> <li>4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;</li> <li>5. Post-seeding drench, transplant-water drench, or hill drench;</li> <li>6. Subsurface side-dress on both sides of each row. ADMIRE PRO SYSTEMIC PROTECTANT must be incorporated into root-zone.</li> </ol> <p><b>Brassica Leafy Vegetables – Soil Application Restrictions</b><br/>           Pre-Harvest Interval (PHI): <b>21 days</b><br/>           Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season: <b>10.5 fluid ounces/Acre</b> (0.38 lb AI/Acre)</p> <p><sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.</p> |                  |                           |   |
|   | <p><b>BRASSICA (COLE) LEAFY VEGETABLES <sup>1/</sup> – FOLIAR</b><br/> <b>Crops of Crop Group 5 including:</b> Broccoli, Broccoli raab (<i>rapini</i>), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (<i>gai lan</i>) broccoli, Chinese (<i>bok choy</i>) cabbage, Chinese (<i>napa</i>) cabbage, Chinese mustard (<i>gai choy</i>) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens</p> <table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Aphids<br/>Flea beetles<br/>Leafhoppers<br/>Whiteflies</td><td>1.3</td></tr> </table> <p><b>Brassica (Cole) Leafy Vegetables – Foliar Application Restrictions</b><br/>           Pre-Harvest Interval (PHI): <b>7 days</b><br/>           Minimum interval between applications: <b>5 days</b><br/>           Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season: <b>6.5 fluid ounces/Acre</b> (0.23 lb AI/Acre)</p> <p><sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.</p>  | Pests Controlled | Rate<br>fluid ounces/Acre | Aphids<br>Flea beetles<br>Leafhoppers<br>Whiteflies |
| Pests Controlled                                    | Rate<br>fluid ounces/Acre  |                  |                           |   |
| Aphids<br>Flea beetles<br>Leafhoppers<br>Whiteflies | 1.3  |                  |                           |   |
| 4.375"  |  |                  |                           |   |







← 3.875" →

**5.5"**

**5.0"**

- ### Legume Vegetables – Soil Application Restrictions

Pre-Harvest Interval (PHI): **21 days**

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.

**Bean** (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)

**Bean** (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

**Other Beans and Peas** [Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean]

### Legume Vegetables – Foliar Application Restrictions

**Pre-Harvest Interval (PHI): 7 days**

Minimum interval between applications: **7 days**

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.

|   |  |                                   |
|---|--|-----------------------------------|
| 3.875"  |  |                                   |
| <p><b>ROOT VEGETABLES – SOIL <sup>1/</sup></b></p> <p><b>Crops of Crop Subgroup 1B except Sugarbeet plus Kava including:</b> Beet (garden)<sup>2/</sup>, Burdock (edible)<sup>2/</sup>, Carrot <sup>2/</sup>, Celeriac<sup>2/</sup>, Chervil (turnip-rooted)<sup>2/</sup>, Chicory<sup>2/</sup>, Ginseng, Horseradish, Kava<sup>2/</sup>, Parsley (turnip-rooted), Parsnip <sup>2/</sup>, Radish <sup>2/</sup>, Oriental radish (diakon)<sup>2/</sup>, Rutabaga<sup>2/</sup>, Salsify (oyster plant), Salsify (black)<sup>2/</sup>, Salsify (Spanish), Skirret and Turnip<sup>2/</sup>.</p> |  |                                   |
| <b>Pests Controlled</b>   | <b>Rate<br/>fluid ounces/1000 row-feet</b> | <b>Rate<br/>fluid ounces/Acre</b> |
| Aphids<br>Flea beetles<br>Leafhoppers<br>Thrips (foliage feeding thrips only)<br>Whiteflies   | 0.31 – 0.74                                | 4.4 – 10.5                        |
| <p><b>Root Vegetables – Soil Applications</b></p> <p>Apply specified dosage in one of the following methods:</p> <ol style="list-style-type: none"> <li>1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;</li> <li>2. In-furrow spray (rate specified per 1000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting;</li> <li>3. In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a bedding operation 14 or fewer days before planting.</li> </ol>                   |  |                                   |
| <p><b>Root Vegetables – Soil Application Restrictions</b></p> <p>Pre-Harvest Interval (PHI): <b>21 days</b></p> <p>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season: <b>10.5 fluid ounces/Acre</b> (0.38 lb AI/Acre)</p> <p>Maximum ADMIRE PRO SYSTEMIC PROTECTANT applications per crop season: <b>1</b></p>   |  |                                   |
| <p><b>Root Vegetables – Soil Application Note</b></p> <p>The rate applied affects the length of control. Use higher rate within the specified rate range where infestations occur later in crop development, or where pest pressure is continuous. ADMIRE PRO SYSTEMIC PROTECTANT rates less than 0.31 fluid ounces/1000 row-feet will not provide adequate residual pest control. ADMIRE PRO SYSTEMIC PROTECTANT treated crops grown on very high organic matter soils (muck) may also require additional pest management control.</p>   |  |                                   |
| <p><sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.</p> <p><sup>2/</sup> Tops or greens from these crops may be utilized for food or feed.</p>  |  |                                   |
| 5.5"  |  |                                   |
| 5.0"  |  |                                   |
| 4.375"  |  |                                   |





| 3.875"  |                           |                  |                           |   |     |
|---|---------------------------|------------------|---------------------------|---|-----|
| <p><b>ROOT, TUBEROUS and CORM VEGETABLES <sup>1/</sup> – FOLIAR</b></p> <p><b>Crops of Crop Group 1C (except sugarbeet) plus Kava including:</b> Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Beet (garden) <sup>2/</sup>, Burdock (edible) <sup>2/</sup>, Canna (edible, Queensland arrowroot), Carrot <sup>2/</sup>, Cassava (bitter &amp; sweet) <sup>2/</sup>, Celeriac <sup>2/</sup>, Chayote (root), Chervil (turnip-rooted) <sup>2/</sup>, Chicory <sup>2/</sup>, Chufa, Dasheen (taro) <sup>2/</sup>, Ginger, Ginseng, Horseradish, Kava <sup>2/,3/</sup>, Leren, Parsley (turnip-rooted), Parsnip <sup>2/</sup>, Radish <sup>2/</sup>, Oriental radish (diakon) <sup>2/</sup>, Rutabaga <sup>2/</sup>, Salsify (black) <sup>2/</sup>, Salsify (oyster plant), Salsify (Spanish), Skirret, Sweetpotato <sup>2/</sup>, Tanier (cocoyam) <sup>2/</sup>, Tumeric, Turnip <sup>2/</sup>, Yam bean (jicama, manioc pea), Yam (true) <sup>2/</sup>.</p> <p><b>(For applications on Potato see Field Crops section)</b></p> <table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Aphids<br/>Flea beetles<br/>Leafhoppers<br/>Whiteflies</td><td>1.2</td></tr> </table> <p><b>Root, Tuberous and Corm Vegetables – Foliar Application Restrictions</b></p> <p>Pre-Harvest Interval (PHI): <b>7 days</b></p> <p>Minimum interval between applications: <b>5 days</b></p> <p>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season on Radish: <b>1.2 fluid ounces/Acre</b> (0.044 lb AI/A);</p> <p>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season on all other crops: <b>3.7 fluid ounces/Acre</b> (0.13 lb AI/A).</p> <p>Maximum ADMIRE PRO SYSTEMIC PROTECTANT applications per crop season: <b>1</b> on Radish;<br/><b>3</b> on other crops.</p> <p><sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.</p> <p><sup>2/</sup> Tops or greens from these crops may be utilized for food or feed.</p> <p><sup>3/</sup> Use not permitted in California unless otherwise directed by state specific 24(c) labeling.</p> |                           | Pests Controlled | Rate<br>fluid ounces/Acre | Aphids<br>Flea beetles<br>Leafhoppers<br>Whiteflies | 1.2 |
| Pests Controlled  | Rate<br>fluid ounces/Acre |                  |                           |   |     |
| Aphids<br>Flea beetles<br>Leafhoppers<br>Whiteflies   | 1.2                       |                  |                           |   |     |
| 5.5"  |                           |                  |                           |   |     |
| 5.0"  |                           |                  |                           |   |     |
| 4.375"  |                           |                  |                           |   |     |

**5.0"**

← 3.875" →

5

| Pests Controlled |            | Rate<br>fluid ounces/Acre |
|------------------|------------|---------------------------|
| Aphids           | Whiteflies | 10.5 – 14.0               |

**Strawberry – Soil (Annual and Perennial) – Applications**

Apply specified dosage in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening;
2. As a plant material or plant hole treatment just prior to, or during transplanting.
3. As a band spray over-the-row in a minimum of 20 gallons of water per acre, followed immediately by overhead irrigation to incorporate product into root-zone. Do not use plastic or other mulches that limit movement of ADMIRE PRO SYSTEMIC PROTECTANT into root zone.

**Strawberry – Soil (Annual and Perennial) – Application Restrictions**

Pre-Harvest Interval (PHI): **14 days**

Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per crop season: **14.0 fluid ounces/Acre** (0.5 lb AI/Acre)

Do not apply immediately prior to bud opening or during bloom or when bees are foraging.

**Strawberry – Soil (Annual and Perennial) – Application Note**

The rate applied affects the length of control. Use higher rate within the specified rate range where infestations may occur later in crop development or where pest pressure is continuous.

<sup>1/</sup> Do not use both soil application methods on the same crop in the same season.

**0**

| Pests Controlled  | Rate<br>fluid ounces/Acre |
|---|---------------------------|
| White grub complex<br>(grubs of Asiatic garden beetle, European and<br>Masked chafer, Japanese beetle, Oriental beetle) | 7.0 – 10.5                |

**Strawberry – Soil (Perennial, Post-Harvest) Applications**

Apply a single application post harvest to coincide with renovation of strawberry fields and during active egg-laying period of beetles. Apply specified dosage in one of the following methods:

1. As a ground spray via boom or backpack sprayer in a minimum of 20 gallons of water per acre;
2. As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth should be equivalent to the width of the anticipated fruiting bed;
3. As a chemigation application with 600 to 1000 gallons of water followed by 0.1 to 0.25 inches irrigation.

33



3.875"

**SUGAR BEET – SOIL <sup>1/</sup> (continued)**

**Sugar beet – Soil Application**

Apply specified dosage in the following method:

1. Apply specified dosage in sufficient carrier volume to insure uniform application. Apply directly below each seed furrow either during the bedding operation immediately prior to planting or at the time of planting.

The low rate may be applied to aid establishment of stands in whitefly areas, or for early season control of the other pests listed.

**Sugar beet – Soil Application Restrictions**

Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: **5.2 fluid ounces/Acre** (0.18 lb AI/Acre)

**Sugar Beet – Soil Application Note**

The low rate may be applied to aid establishment of stands in whitefly areas, or for early season control of the other pests listed.

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state specific 24(c) labeling.

**ADMIRE PRO SYSTEMIC PROTECTANT  
CONVERSION CHART FOR LINEAR APPLICATION**

| RATE<br>fluid ounces per Acre | RATE<br>fluid ounces/1000 row-feet               |      |      |      |      |      |      |      |      |      |
|-------------------------------|--|------|------|------|------|------|------|------|------|------|
|                               | Based on <i>average</i> row spacing (in inches): |      |      |      |      |      |      |      |      |      |
|                               | 10   | 15   | 20   | 25   | 30   | 34   | 36   | 38   | 40   | 45   |
| 5.0                           | 0.10   | 0.14 | 0.19 | 0.24 | 0.29 | 0.33 | 0.34 | 0.36 | 0.38 | 0.43 |
| 5.5                           | 0.11   | 0.16 | 0.21 | 0.26 | 0.32 | 0.36 | 0.38 | 0.40 | 0.42 | 0.47 |
| 6.0                           | 0.11   | 0.17 | 0.23 | 0.29 | 0.34 | 0.39 | 0.41 | 0.44 | 0.46 | 0.52 |
| 6.5                           | 0.12   | 0.19 | 0.25 | 0.31 | 0.37 | 0.42 | 0.45 | 0.47 | 0.50 | 0.56 |
| 7.0                           | 0.13   | 0.20 | 0.27 | 0.33 | 0.40 | 0.46 | 0.48 | 0.51 | 0.54 | 0.60 |
| 7.5                           | 0.14   | 0.22 | 0.29 | 0.36 | 0.43 | 0.49 | 0.52 | 0.55 | 0.57 | 0.65 |
| 8.0                           | 0.15   | 0.23 | 0.31 | 0.38 | 0.46 | 0.52 | 0.55 | 0.58 | 0.61 | 0.69 |
| 8.5                           | 0.16   | 0.24 | 0.33 | 0.41 | 0.49 | 0.55 | 0.59 | 0.62 | 0.65 | 0.73 |
| 9.0                           | 0.17   | 0.26 | 0.34 | 0.43 | 0.52 | 0.59 | 0.62 | 0.65 | 0.69 | 0.77 |
| 9.5                           | 0.18   | 0.27 | 0.36 | 0.45 | 0.55 | 0.62 | 0.65 | 0.69 | 0.73 | 0.82 |
| 10.0                          | 0.19   | 0.29 | 0.38 | 0.48 | 0.57 | 0.65 | 0.69 | 0.73 | 0.77 | 0.86 |
| 10.5                          | 0.20   | 0.30 | 0.40 | 0.50 | 0.60 | 0.68 | 0.72 | 0.76 | 0.80 | 0.90 |

4.375"

3.875"

**ADMIRE PRO SYSTEMIC PROTECTANT  
CONVERSION CHART FOR LINEAR APPLICATION (continued)**

| RATE<br>fluid ounces per Acre | RATE<br>fluid ounces/1000 row-feet               |      |      |      |      |      |      |      |      |      |
|-------------------------------|--|------|------|------|------|------|------|------|------|------|
|                               | Based on <i>average</i> row spacing (in inches): |      |      |      |      |      |      |      |      |      |
|                               | 10   | 15   | 20   | 25   | 30   | 34   | 36   | 38   | 40   | 45   |
| 11.0                          | 0.21   | 0.32 | 0.42 | 0.53 | 0.63 | 0.72 | 0.76 | 0.80 | 0.84 | 0.95 |
| 11.5                          | 0.22   | 0.33 | 0.44 | 0.55 | 0.66 | 0.75 | 0.79 | 0.84 | 0.88 | 0.99 |
| 12.0                          | 0.23   | 0.34 | 0.46 | 0.57 | 0.69 | 0.78 | 0.83 | 0.87 | 0.92 | 1.03 |
| 12.5                          | 0.24   | 0.36 | 0.48 | 0.60 | 0.72 | 0.81 | 0.86 | 0.91 | 0.96 | 1.08 |
| 13.0                          | 0.25   | 0.37 | 0.50 | 0.62 | 0.75 | 0.85 | 0.90 | 0.95 | 0.99 | 1.12 |
| 13.5                          | 0.26   | 0.39 | 0.52 | 0.65 | 0.77 | 0.88 | 0.93 | 0.98 | 1.03 | 1.16 |
| 14.0                          | 0.27   | 0.40 | 0.54 | 0.67 | 0.80 | 0.91 | 0.96 | 1.02 | 1.07 | 1.21 |

**Important Note:** The ADMIRE PRO SYSTEMIC PROTECTANT rate applied affects the length of control and to a considerable extent, the degree of control or effect. Row-spacing X ADMIRE PRO SYSTEMIC PROTECTANT rate combinations in shaded blocks may not provide adequate residual pest control and are not recommended for long-term, residual control. Use higher labeled rates where infestations may occur later in crop development or where pest pressure is continuous. Bayer CropScience offers no warranty for use of ADMIRE PRO SYSTEMIC PROTECTANT at rates below 0.31 fluid ounces/1000 row-feet.

**TREE, BUSH and VINE CROPS  
Application Rates – ADMIRE PRO Systemic Protectant**

**BANANA and PLANTAIN – SOIL**

| Pests Controlled   | Rate<br>fluid ounces/Acre |
|--|---------------------------|
| Aphids      Leafhoppers  | 7.0 – 14.0                |
| <b>Pest Suppressed</b>   |                           |
| Scales   | 7.0 – 14.0                |
| <b>Banana and Plantain – Soil Application</b>  |                           |
| Apply specified dosage of ADMIRE PRO SYSTEMIC PROTECTANT in the following method:                          |                           |
| 1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. |                           |
| <b>Banana and Plantain – Soil Application Restrictions</b>   |                           |
| Pre-Harvest Interval (PHI): 0 day  |                           |
| Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>14.0 fluid ounces/Acre</b> (0.5 lb AI/A)       |                           |

4.375"









3.875"

#### CANEBERRY – FOLIAR

**Crops of Crop Subgroup 13A including:**

**Blackberry** (*Rubus eubatus*, including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these)

**Raspberry** (black and red, *Rubus occidentalis*, *Rubus strigosus*, *Rubus idaeus*)

| Pests Controlled  | Rate<br>fluid ounces/Acre |
|---|---------------------------|
| Aphids<br>Leafhoppers<br>Thrips (foliage feeding thrips, only)  | 2.8                       |
| <b>Caneberry – Foliar Application Restrictions</b><br>Pre-Harvest Interval (PHI): <b>3 days</b><br>Minimum interval between applications: <b>7 days</b><br>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>8.4 fluid ounces/Acre</b><br>(0.3 lb AI/Acre)<br>Do not apply pre-bloom or during bloom or when bees are foraging. |                           |

#### CITRUS (Containerized) – SOIL

**Crops of Crop Group 10 Including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, and other cultivars and/or hybrids of these.

| Pests Controlled  | Rate<br>mL / "citra pot" (0.1 ft³ container media)                          |
|---|---|
| Aphids<br>Asian citrus psyllid<br>Blackfly<br>Citrus leafminer<br>Leafhoppers/Sharpshooters | Mealybugs<br>Scales<br>Whiteflies<br>Citrus root weevil<br>(larval complex) |
|   | 0.33 – 0.50   |
| <b>Pest Suppressed</b>  |   |
| Citrus thrips (foliage feeding thrips only)   | 0.50  |

(continued)

4.375"



|  |  | 3.875"  |                  |      |                           |   |   |            |  |  |
|--|--|---|------------------|------|---------------------------|---|---|------------|--|--|
| 5.5"   | <b>CITRUS (Field) – SOIL</b>   |   |                  | 5.0" |                           |   |   |            |  |  |
|  | <b>Crops of Crop Group 10 Including:</b> Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, and other cultivars and/or hybrids of these.                            |   |                  |      |                           |   |   |            |  |  |
|  | <table><tr><th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr><tr><td>Aphids<br/>Asian citrus psyllid<br/>Blackfly<br/>Citrus leafminer<br/>Leafhoppers/Sharpshooters</td><td>Mealybugs<br/>Scales<br/>Termites (FL only)<br/>Whiteflies</td></tr><tr><td colspan="2">7.0 – 14.0</td></tr></table> |   | Pests Controlled |      | Rate<br>fluid ounces/Acre | Aphids<br>Asian citrus psyllid<br>Blackfly<br>Citrus leafminer<br>Leafhoppers/Sharpshooters | Mealybugs<br>Scales<br>Termites (FL only)<br>Whiteflies | 7.0 – 14.0 |  |  |
|  | Pests Controlled   | Rate<br>fluid ounces/Acre                               |                  |      |                           |   |   |            |  |  |
|  | Aphids<br>Asian citrus psyllid<br>Blackfly<br>Citrus leafminer<br>Leafhoppers/Sharpshooters  | Mealybugs<br>Scales<br>Termites (FL only)<br>Whiteflies |                  |      |                           |   |   |            |  |  |
| 7.0 – 14.0   |  |   |                  |      |                           |   |   |            |  |  |
| <b>Pests / Diseases Suppressed</b>   |  |   |                  |      |                           |   |   |            |  |  |
| Citrus nematode<br>Symptoms of:<br>Citrus tristeza virus (CTV) through vector control<br>Citrus yellows<br>Thrips (foliage feeding thrips only)  |  | 14.0  |                  |      |                           |   |   |            |  |  |
| <b>Citrus (Field) – Soil Applications</b><br>Apply specified dosage in one of the following methods:<br>1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications of ADMIRE PRO SYSTEMIC PROTECTANT. Chemigation application can be made separate to normal irrigation but followed by 10 to 20 minutes of additional watering to move ADMIRE PRO SYSTEMIC PROTECTANT into root-zone. Allow 24 hours before initiating subsequent irrigations;<br>2. Band spray soil surface on both sides of the tree. Overlap bands at the tree base to create a continuous band within the drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root-zone. This method is suitable for very coarse soils with 0.75% organic matter or less;<br>3. Drench to base of tree not exceeding one-quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. For Florida citrus, only use this treatment method for trees up to 8 feet tall;<br>4. For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk.<br>5. For suppression of citrus nematode, apply specified dosage through low-pressure chemigation or soil surface band spray only, ensuring complete coverage of the root system and utilizing application directions stated above for the respective application method. Repeated and regular use of ADMIRE PRO SYSTEMIC PROTECTANT over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response. |  |   |                  |      |                           |   |   |            |  |  |
| <b>Citrus (Field) – Soil Applications Restrictions</b><br>Pre-Harvest Interval (PHI): <b>0 day</b><br>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>14.0 fluid ounces/Acre</b> (0.5 lb AI/Acre)  |  |   |                  |      |                           |   |   |            |  |  |
|  |  | 42  |                  |      |                           |   |   |            |  |  |
|  |  | 4.375"  |                  |      |                           |   |   |            |  |  |





3.875"

A vertical double-headed arrow with the text "5.5''" centered next to it, indicating a height of 5.5 inches.

**5.0"**

|  |
|--|
| <p><b>Cranberry – Soil Application Restrictions</b></p> <p>Pre-Harvest Interval (PHI): <b>30 days</b></p> <p>Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>14.0 fluid ounces/Acre</b> (0.5 lb AI/Acre)</p> <p>Do not apply immediately pre-bloom or during bloom or when bees are foraging</p> |
|--|

Best control may be achieved when application is made post-bloom immediately after bees are removed. Applications should target early instar larvae.

ADMIRE PRO SYSTEMIC PROTECTANT has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of the ADMIRE PRO SYSTEMIC PROTECTANT and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.

**Including:** American bunch grape, Muscadine grape and Vinifera grape

### Grape – Soil Applications

Apply specified dosage in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation;
3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation.
4. For suppression of nematodes, apply 14 fluid ounces in a single application or two 7 fluid ounce applications on a 30 to 45 day interval. Apply only by 1) chemigation into root-zone through above ground low-pressure drip, trickle, micro-sprinkler or equivalent equipment; or 2) French plow technique, followed immediately by sufficient irrigation to move the product into the entire root-zone of the plant. Repeated and regular use of ADMIRE PRO SYSTEMIC PROTECTANT over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.

(continued)

← 4.375" →











3.875"

A vertical double-headed arrow with the text "5.5''" centered next to it, indicating a height of 5.5 inches.

A vertical dimension line with arrows at both ends, indicating a height of 5.0 inches. The text "5.0" is centered next to the line.


**Crops Of Crop Group 12 Including:** Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

| <b>Pest Controlled</b>  | <b>Rate<br/>fluid ounces/10 gallons root-dip solution</b> |
|---|---|
| Black peach aphid (infesting roots)   | 0.87 (26 mLs)   |
| <b>Stone Fruit – Pre-plant Root-dip Applications</b><br>Mix ADMIRE PRO SYSTEMIC PROTECTANT at 0.87 fluid ounces (26 mLs) per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in the ADMIRE PRO SYSTEMIC PROTECTANT solution for up to 5 minutes. Allow solution to dry on roots and transplant trees as soon as possible following treatment. |   |



| 3.875"  |   |                  |                           |   |           |  |     |
|---|---|------------------|---------------------------|---|-----------|--|-----|
| 5.5"  | <p><b>TREE NUTS – FOLIAR</b><br/> <b>Crops of Crop Group 14 except Almond including:</b> Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut [black and English]</p> <table> <tr> <th>Pests Controlled</th><th>Rate<br/>fluid ounces/Acre</th></tr> <tr> <td>Aphids (except Black pecan aphid)<br/>Leafhoppers/Sharpshooters<br/><i>Phylloxera</i> spp. (leaf infestations)<br/>Spittlebugs<br/>Whiteflies</td><td>1.2 – 2.4</td></tr> <tr> <td>Black pecan aphid<br/>Mealybugs<br/>San Jose scale</td><td>2.8</td></tr> </table> <p><b>Tree Nuts – Foliar Applications</b><br/> Applications for control of San Jose scale should be timed according to crawler stage, treating each successive generation. Two applications on a 10 to 14-day interval may be required to achieve control.</p> <p><b>Tree Nuts – Foliar Application Restrictions</b><br/> Pre-Harvest Interval (PHI): <b>7 days</b><br/> Minimum interval between applications: <b>6 days</b><br/> Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per year: <b>10.0 fluid ounces/Acre</b><br/> (0.36 lb AI/A)<br/> Minimum application volume (water): 50 GPA – ground application, 25 GPA – aerial application<br/> Do not apply pre-bloom or during bloom or when bees are foraging.</p> | Pests Controlled | Rate<br>fluid ounces/Acre | Aphids (except Black pecan aphid)<br>Leafhoppers/Sharpshooters<br><i>Phylloxera</i> spp. (leaf infestations)<br>Spittlebugs<br>Whiteflies | 1.2 – 2.4 | Black pecan aphid<br>Mealybugs<br>San Jose scale | 2.8 |
| Pests Controlled  | Rate<br>fluid ounces/Acre   |                  |                           |   |           |  |     |
| Aphids (except Black pecan aphid)<br>Leafhoppers/Sharpshooters<br><i>Phylloxera</i> spp. (leaf infestations)<br>Spittlebugs<br>Whiteflies | 1.2 – 2.4   |                  |                           |   |           |  |     |
| Black pecan aphid<br>Mealybugs<br>San Jose scale  | 2.8   |                  |                           |   |           |  |     |
| 5.0"  |   |                  |                           |   |           |  |     |
| 4.375"  |   |                  |                           |   |           |  |     |









| 3.875"   |  |                 |  |                        |   |                         |  |                                       |   |
|--|--|-----------------|--|------------------------|---|-------------------------|--|---------------------------------------|---|
| <p><b>POPLAR / COTTONWOOD – CUTTINGS/WHIPS SOAK <sup>1/</sup></b><br/> (includes members of the genus <i>Populus</i> grown for pulp or timber)</p> <table> <tr> <th>Pest Controlled</th><th>Cuttings/Whips Soaking Solution<br/>fluid ounces ADMIRE PRO SYSTEMIC<br/>PROTECTANT Needed per 100 gallons</th></tr> <tr> <td>Cottonwood leaf beetle</td><td>5.8 – 11.6 (unhydrated cuttings/whips)<br/>11.6 – 17.5 (partially hydrated cuttings/whips)</td></tr> <tr> <td colspan="2"><b>Pests Suppressed</b></td></tr> <tr> <td>Aphids<br/><i>Phylloxera popularia</i></td><td>5.8 – 11.6 (unhydrated cuttings/whips)<br/>11.6 – 17.5 (partially hydrated cuttings/whips)</td></tr> </table> <p><b>Poplar / Cottonwood – Cuttings/Whips Applications</b><br/> Moisture content of cuttings/whips prior to application, the solution concentration and the length of soaking interval interact to affect the amount of product absorbed into plant material. For a constant soaking interval of 24 hours, drier cuttings/whips absorb a higher quantity of solution and require a lower concentration. Conversely, more hydrated cuttings/whips absorb less solution and require a higher concentration. Soaking of cuttings/whips should occur in a covered container in absence of UV light. Not all <i>Populus</i> sp. clones/varieties/hybrids have been tested for crop safety. Without specific knowledge about a particular <i>Populus</i> sp. clone/variety/hybrid, Bayer CropScience recommends that small numbers of cuttings/whips of each be treated and evaluated prior to commercial use.</p> <p><u>Apply ADMIRE PRO SYSTEMIC PROTECTANT in one of the following cuttings/whips soaking methods:</u></p> <ol style="list-style-type: none"> <li>1. For freshly cut (unhydrated) cuttings/whips, soak plant material in specified solution concentration for 24 hours prior to cold storage. After removal from cold storage, plant as needed.</li> <li>2. For previously hydrated cuttings/whips removed from cold storage, allow plant material to reach room temperature and soak in specified solution concentration for 24 hours prior to planting.</li> </ol> <p>Proper care must be taken in disposal of any residual soaking solution. Solution may be applied to existing trees or other registered crops as long as all product label precautions and restrictions are observed.</p> <p><b>Poplar / Cottonwood – Cuttings/Whips Applications Restrictions</b><br/> Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed at-plant per year: <b>14.0 fluid ounces/Acre</b> (0.5 lb AI/Acre)</p> <p><sup>1/</sup> Use not permitted in California unless otherwise directed by state specific 24(c) labeling.</p> |  | Pest Controlled | Cuttings/Whips Soaking Solution<br>fluid ounces ADMIRE PRO SYSTEMIC<br>PROTECTANT Needed per 100 gallons | Cottonwood leaf beetle | 5.8 – 11.6 (unhydrated cuttings/whips)<br>11.6 – 17.5 (partially hydrated cuttings/whips) | <b>Pests Suppressed</b> |  | Aphids<br><i>Phylloxera popularia</i> | 5.8 – 11.6 (unhydrated cuttings/whips)<br>11.6 – 17.5 (partially hydrated cuttings/whips) |
| Pest Controlled  | Cuttings/Whips Soaking Solution<br>fluid ounces ADMIRE PRO SYSTEMIC<br>PROTECTANT Needed per 100 gallons |                 |  |                        |   |                         |  |                                       |   |
| Cottonwood leaf beetle   | 5.8 – 11.6 (unhydrated cuttings/whips)<br>11.6 – 17.5 (partially hydrated cuttings/whips)                |                 |  |                        |   |                         |  |                                       |   |
| <b>Pests Suppressed</b>  |  |                 |  |                        |   |                         |  |                                       |   |
| Aphids<br><i>Phylloxera popularia</i>  | 5.8 – 11.6 (unhydrated cuttings/whips)<br>11.6 – 17.5 (partially hydrated cuttings/whips)                |                 |  |                        |   |                         |  |                                       |   |
| 5.5"   |  |                 |  |                        |   |                         |  |                                       |   |
| 5.0"   |  |                 |  |                        |   |                         |  |                                       |   |
| 4.375"   |  |                 |  |                        |   |                         |  |                                       |   |

$\longleftrightarrow 3.875'' \longrightarrow$

(includes members of the genus *Populus* grown for pulp or timber)

**5.5"**


**5.0"**


**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.





59

.8125"

4.0"

.5625"

NET CONTENTS: 1 GAL. 12 OZ. (140 FL. OZ.)

5.25"



5.5"

5.375"

## ADMIRE® PRO Systemic Protectant

For uses in pest management and maintenance of plant health.

ACTIVE INGREDIENT: Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine . . . . . 42.8%  
OTHER INGREDIENTS: . . . . . 57.2%  
Contains 4.6 pounds of active ingredient per gallon or 550 grams AI/liter. TOTAL: 100.0%  
SHAKE WELL BEFORE USING

EPA Reg. No. 264-827

**STOP - Read the label before use**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call  
24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER  
(1-866-992-2937)

See Panel for First Aid Instructions and Booklet for Complete  
Precautionary Statements and Directions for Use.

### FIRST AID

|                        |   |
|------------------------|---|
| IF SWALLOWED           | <ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul> |
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>  |
| IF INHALED             | <ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>   |
| IF IN EYES             | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>  |

In case of emergency call toll free the Bayer CropScience  
Emergency Response Telephone No. 1-800-334-7577.  
Have a product container or label with you when calling a  
poison control center or doctor, or going for treatment.

### GROUP 4A INSECTICIDE

**Note To Physician:** No specific antidote is available. Treat the patient symptomatically.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response telephone number is 1-800-334-7577.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
ADMIRE is a registered trademark of Bayer.  
©2019 Bayer CropScience

US6138118B8 190917B 10/19